

## Zero-Based Finite Sequences

Tetsuya Tsunetou  
Kyushu University

Grzegorz Bancerek  
Shinshu University  
Nagano

Yatsuka Nakamura  
Shinshu University  
Nagano

MML Identifier: AFINSQ\_1.

WWW: [http://mizar.org/JFM/Vol13/afinsq\\_1.html](http://mizar.org/JFM/Vol13/afinsq_1.html)

The articles [14], [10], [16], [15], [17], [8], [3], [4], [5], [13], [2], [9], [12], [1], [6], [11], and [7] provide the notation and terminology for this paper.

We follow the rules:  $k, n$  denote natural numbers,  $x, y, z, y_1, y_2, X$  denote sets, and  $f$  denotes a function.

One can prove the following propositions:

- (1)  $n \in n + 1$ .
- (2) If  $k \leq n$ , then  $k = k \cap n$ .
- (3) If  $k = k \cap n$ , then  $k \leq n$ .
- (4)  $n \cup \{n\} = n + 1$ .
- (5)  $\text{Seg } n \subseteq n + 1$ .
- (6)  $n + 1 = \{0\} \cup \text{Seg } n$ .
- (7) For every function  $r$  holds  $r$  is finite and transfinite sequence-like iff there exists  $n$  such that  $\text{dom } r = n$ .

Let us observe that there exists a function which is finite and transfinite sequence-like.

A finite 0-sequence is a finite transfinite sequence.

In the sequel  $p, q, r$  denote finite 0-sequences.

Observe that every set which is natural is also finite. Let us consider  $p$ . One can check that  $\text{dom } p$  is natural.

Let us consider  $p$ . Then  $\overline{\overline{p}}$  is a natural number and it can be characterized by the condition:

(Def. 1)  $\overline{\overline{p}} = \text{dom } p$ .

We introduce  $\text{len } p$  as a synonym of  $\overline{\overline{p}}$ .

Let us consider  $p$ . Then  $\text{dom } p$  is a subset of  $\mathbb{N}$ .

We now state the proposition

- (8) If there exists  $k$  such that  $\text{dom } f \subseteq k$ , then there exists  $p$  such that  $f \subseteq p$ .

In this article we present several logical schemes. The scheme *XSeqEx* deals with a natural number  $\mathcal{A}$  and a binary predicate  $\mathcal{P}$ , and states that:

There exists  $p$  such that  $\text{dom } p = \mathcal{A}$  and for every  $k$  such that  $k \in \mathcal{A}$  holds  $\mathcal{P}[k, p(k)]$

provided the following requirements are met:

- For all  $k, y_1, y_2$  such that  $k \in \mathcal{A}$  and  $\mathcal{P}[k, y_1]$  and  $\mathcal{P}[k, y_2]$  holds  $y_1 = y_2$ , and
- For every  $k$  such that  $k \in \mathcal{A}$  there exists  $x$  such that  $\mathcal{P}[k, x]$ .

The scheme *XSeqLambda* deals with a natural number  $\mathcal{A}$  and a unary functor  $\mathcal{F}$  yielding a set, and states that:

There exists a finite 0-sequence  $p$  such that  $\text{len } p = \mathcal{A}$  and for every  $k$  such that  $k \in \mathcal{A}$  holds  $p(k) = \mathcal{F}(k)$

for all values of the parameters.

Next we state several propositions:

- (9) If  $z \in p$ , then there exists  $k$  such that  $k \in \text{dom } p$  and  $z = \langle k, p(k) \rangle$ .
- (10) If  $\text{dom } p = \text{dom } q$  and for every  $k$  such that  $k \in \text{dom } p$  holds  $p(k) = q(k)$ , then  $p = q$ .
- (11) If  $\text{len } p = \text{len } q$  and for every  $k$  such that  $k < \text{len } p$  holds  $p(k) = q(k)$ , then  $p = q$ .
- (12)  $p \upharpoonright n$  is a finite 0-sequence.
- (13) If  $\text{rng } p \subseteq \text{dom } f$ , then  $f \cdot p$  is a finite 0-sequence.
- (14) If  $k < \text{len } p$  and  $q = p \upharpoonright k$ , then  $\text{len } q = k$  and  $\text{dom } q = k$ .

Let  $D$  be a set. Observe that there exists a transfinite sequence of elements of  $D$  which is finite.

Let  $D$  be a set. A finite 0-sequence of  $D$  is a finite transfinite sequence of elements of  $D$ .

The following proposition is true

- (15) For every set  $D$  holds every finite 0-sequence of  $D$  is a partial function from  $\mathbb{N}$  to  $D$ .

One can check that  $\emptyset$  is transfinite sequence-like.

Let  $D$  be a set. One can verify that there exists a partial function from  $\mathbb{N}$  to  $D$  which is finite and transfinite sequence-like.

In the sequel  $D$  is a set.

One can prove the following propositions:

- (16) For every finite 0-sequence  $p$  of  $D$  holds  $p \upharpoonright k$  is a finite 0-sequence of  $D$ .
- (17) For every non empty set  $D$  there exists a finite 0-sequence  $p$  of  $D$  such that  $\text{len } p = k$ .

Let us note that there exists a finite 0-sequence which is empty.

Next we state two propositions:

- (18)  $\text{len } p = 0$  iff  $p = \emptyset$ .
- (19) For every set  $D$  holds  $\emptyset$  is a finite 0-sequence of  $D$ .

Let  $D$  be a set. Note that there exists a finite 0-sequence of  $D$  which is empty.

Let us consider  $x$ . The functor  $\langle \_0 x \rangle$  yields a set and is defined by:

(Def. 2)  $\langle \_0 x \rangle = \{ \langle 0, x \rangle \}$ .

Let  $D$  be a set. The functor  $\langle \_ \rangle_D$  yields an empty finite 0-sequence of  $D$  and is defined as follows:

(Def. 3)  $\langle \_ \rangle_D = \emptyset$ .

Let us consider  $p, q$ . Note that  $p \hat{\ } q$  is finite. Then  $p \hat{\ } q$  can be characterized by the condition:

(Def. 4)  $\text{dom}(p \hat{\ } q) = \text{len } p + \text{len } q$  and for every  $k$  such that  $k \in \text{dom } p$  holds  $(p \hat{\ } q)(k) = p(k)$  and for every  $k$  such that  $k \in \text{dom } q$  holds  $(p \hat{\ } q)(\text{len } p + k) = q(k)$ .

Next we state a number of propositions:

- (20)  $\text{len}(p \hat{\ } q) = \text{len } p + \text{len } q$ .

- (21) If  $\text{len } p \leq k$  and  $k < \text{len } p + \text{len } q$ , then  $(p \hat{\ } q)(k) = q(k - \text{len } p)$ .
- (22) If  $\text{len } p \leq k$  and  $k < \text{len}(p \hat{\ } q)$ , then  $(p \hat{\ } q)(k) = q(k - \text{len } p)$ .
- (23) If  $k \in \text{dom}(p \hat{\ } q)$ , then  $k \in \text{dom } p$  or there exists  $n$  such that  $n \in \text{dom } q$  and  $k = \text{len } p + n$ .
- (24) For all transfinite sequences  $p, q$  holds  $\text{dom } p \subseteq \text{dom}(p \hat{\ } q)$ .
- (25) If  $x \in \text{dom } q$ , then there exists  $k$  such that  $k = x$  and  $\text{len } p + k \in \text{dom}(p \hat{\ } q)$ .
- (26) If  $k \in \text{dom } q$ , then  $\text{len } p + k \in \text{dom}(p \hat{\ } q)$ .
- (27)  $\text{rng } p \subseteq \text{rng}(p \hat{\ } q)$ .
- (28)  $\text{rng } q \subseteq \text{rng}(p \hat{\ } q)$ .
- (29)  $\text{rng}(p \hat{\ } q) = \text{rng } p \cup \text{rng } q$ .
- (30)  $(p \hat{\ } q) \hat{\ } r = p \hat{\ } (q \hat{\ } r)$ .
- (31) If  $p \hat{\ } r = q \hat{\ } r$  or  $r \hat{\ } p = r \hat{\ } q$ , then  $p = q$ .
- (32)  $p \hat{\ } \emptyset = p$  and  $\emptyset \hat{\ } p = p$ .
- (33) If  $p \hat{\ } q = \emptyset$ , then  $p = \emptyset$  and  $q = \emptyset$ .

Let  $D$  be a set and let  $p, q$  be finite 0-sequences of  $D$ . Then  $p \hat{\ } q$  is a transfinite sequence of elements of  $D$ .

Let us consider  $x$ . Then  $\langle 0x \rangle$  is a function and it can be characterized by the condition:

(Def. 5)  $\text{dom } \langle 0x \rangle = 1$  and  $\langle 0x \rangle(0) = x$ .

Let us consider  $x$ . Note that  $\langle 0x \rangle$  is function-like and relation-like.

Let us consider  $x$ . Observe that  $\langle 0x \rangle$  is finite and transfinite sequence-like.

We now state the proposition

(34) Suppose  $p \hat{\ } q$  is a finite 0-sequence of  $D$ . Then  $p$  is a finite 0-sequence of  $D$  and  $q$  is a finite 0-sequence of  $D$ .

Let us consider  $x, y$ . The functor  $\langle 0x, y \rangle$  yielding a set is defined by:

(Def. 6)  $\langle 0x, y \rangle = \langle 0x \rangle \hat{\ } \langle 0y \rangle$ .

Let us consider  $z$ . The functor  $\langle 0x, y, z \rangle$  yielding a set is defined as follows:

(Def. 7)  $\langle 0x, y, z \rangle = \langle 0x \rangle \hat{\ } \langle 0y \rangle \hat{\ } \langle 0z \rangle$ .

Let us consider  $x, y$ . One can verify that  $\langle 0x, y \rangle$  is function-like and relation-like. Let us consider  $z$ . Observe that  $\langle 0x, y, z \rangle$  is function-like and relation-like.

Let us consider  $x, y$ . Note that  $\langle 0x, y \rangle$  is finite and transfinite sequence-like. Let us consider  $z$ . Observe that  $\langle 0x, y, z \rangle$  is finite and transfinite sequence-like.

One can prove the following propositions:

- (35)  $\langle 0x \rangle = \{\langle 0, x \rangle\}$ .
- (36)  $p = \langle 0x \rangle$  iff  $\text{dom } p = 1$  and  $\text{rng } p = \{x\}$ .
- (37)  $p = \langle 0x \rangle$  iff  $\text{len } p = 1$  and  $\text{rng } p = \{x\}$ .
- (38)  $p = \langle 0x \rangle$  iff  $\text{len } p = 1$  and  $p(0) = x$ .
- (39)  $(\langle 0x \rangle \hat{\ } p)(0) = x$ .
- (40)  $(p \hat{\ } \langle 0x \rangle)(\text{len } p) = x$ .

- (41)  $\langle 0x, y, z \rangle = \langle 0x \rangle \hat{\ } \langle 0y, z \rangle$  and  $\langle 0x, y, z \rangle = \langle 0x, y \rangle \hat{\ } \langle 0z \rangle$ .
- (42)  $p = \langle 0x, y \rangle$  iff  $\text{len } p = 2$  and  $p(0) = x$  and  $p(1) = y$ .
- (43)  $p = \langle 0x, y, z \rangle$  iff  $\text{len } p = 3$  and  $p(0) = x$  and  $p(1) = y$  and  $p(2) = z$ .
- (44) If  $p \neq \emptyset$ , then there exist  $q, x$  such that  $p = q \hat{\ } \langle 0x \rangle$ .

Let  $D$  be a non empty set and let  $x$  be an element of  $D$ . Then  $\langle 0x \rangle$  is a finite 0-sequence of  $D$ .  
The scheme *IndXSeq* concerns a unary predicate  $\mathcal{P}$ , and states that:

For every  $p$  holds  $\mathcal{P}[p]$

provided the parameters meet the following requirements:

- $\mathcal{P}[\emptyset]$ , and
- For all  $p, x$  such that  $\mathcal{P}[p]$  holds  $\mathcal{P}[p \hat{\ } \langle 0x \rangle]$ .

We now state the proposition

- (45) For all finite 0-sequences  $p, q, r, s$  such that  $p \hat{\ } q = r \hat{\ } s$  and  $\text{len } p \leq \text{len } r$  there exists a finite 0-sequence  $t$  such that  $p \hat{\ } t = r$ .

Let  $D$  be a set. The functor  $D^0$  yields a set and is defined as follows:

(Def. 8)  $x \in D^0$  iff  $x$  is a finite 0-sequence of  $D$ .

Let  $D$  be a set. Note that  $D^0$  is non empty.

The following propositions are true:

- (46)  $x \in D^0$  iff  $x$  is a finite 0-sequence of  $D$ .
- (47)  $\emptyset \in D^0$ .

The scheme *SepXSeq* deals with a non empty set  $\mathcal{A}$  and a unary predicate  $\mathcal{P}$ , and states that:

There exists  $X$  such that for every  $x$  holds  $x \in X$  iff there exists  $p$  such that  $p \in \mathcal{A}^0$   
and  $\mathcal{P}[p]$  and  $x = p$

for all values of the parameters.

Let  $p$  be a finite 0-sequence and let  $i, x$  be sets. One can check that  $p + \cdot (i, x)$  is finite and transfinite sequence-like. We introduce  $\text{Replace}(p, i, x)$  as a synonym of  $p + \cdot (i, x)$ .

The following proposition is true

- (48) Let  $p$  be a finite 0-sequence,  $i$  be a natural number, and  $x$  be a set. Then  $\text{len } \text{Replace}(p, i, x) = \text{len } p$  and if  $i < \text{len } p$ , then  $(\text{Replace}(p, i, x))(i) = x$  and for every natural number  $j$  such that  $j \neq i$  holds  $(\text{Replace}(p, i, x))(j) = p(j)$ .

Let  $D$  be a non empty set, let  $p$  be a finite 0-sequence of  $D$ , let  $i$  be a natural number, and let  $a$  be an element of  $D$ . Then  $\text{Replace}(p, i, a)$  is a finite 0-sequence of  $D$ .

## REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/card\\_1.html](http://mizar.org/JFM/Vol1/card_1.html).
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/nat\\_1.html](http://mizar.org/JFM/Vol1/nat_1.html).
- [3] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [4] Grzegorz Bancerek. Sequences of ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal2.html>.
- [5] Grzegorz Bancerek. Increasing and continuous ordinal sequences. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/ordinal4.html>.
- [6] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finseq\\_1.html](http://mizar.org/JFM/Vol1/finseq_1.html).
- [7] Grzegorz Bancerek and Andrzej Trybulec. Miscellaneous facts about functions. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/funct\\_7.html](http://mizar.org/JFM/Vol8/funct_7.html).

- [8] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_1.html](http://mizar.org/JFM/Vol1/funct_1.html).
- [9] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/partfun1.html>.
- [10] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/zfmisc\\_1.html](http://mizar.org/JFM/Vol1/zfmisc_1.html).
- [11] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/funct\\_4.html](http://mizar.org/JFM/Vol2/funct_4.html).
- [12] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finset\\_1.html](http://mizar.org/JFM/Vol1/finset_1.html).
- [13] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/real\\_1.html](http://mizar.org/JFM/Vol1/real_1.html).
- [14] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [15] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [16] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/subset\\_1.html](http://mizar.org/JFM/Vol1/subset_1.html).
- [17] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relat\\_1.html](http://mizar.org/JFM/Vol1/relat_1.html).

*Received September 28, 2001*

*Published January 2, 2004*

---