

Atlas of Midpoint Algebra

Michał Muzalewski
Warsaw University
Białystok

Summary. This article is a continuation of [5]. We have established a one-to-one correspondence between midpoint algebras and groups with the operator $1/2$. In general we shall say that a given midpoint algebra M and a group V are w -associated iff w is an atlas from M to V . At the beginning of the paper a few facts which rather belong to [4], [?] are proved.

MML Identifier: MIDSP_2.

WWW: http://mizar.org/JFM/Vol3/midsp_2.html

The articles [3], [7], [2], [1], [6], [4], and [5] provide the notation and terminology for this paper.

For simplicity, we adopt the following rules: G denotes a non empty loop structure, x denotes an element of G , M denotes a non empty midpoint algebra structure, p, q, r denote points of M , and w denotes a function from $[\text{the carrier of } M, \text{ the carrier of } M]$ into the carrier of G .

Let us consider G, x . The functor $2x$ yielding an element of G is defined as follows:

(Def. 1) $2x = x + x$.

Let us consider M, G, w . We say that M, G are associated w.r.t. w if and only if:

(Def. 2) $p @ q = r$ iff $w(p, r) = w(r, q)$.

The following proposition is true

(1) If M, G are associated w.r.t. w , then $p @ p = p$.

We use the following convention: S denotes a non empty set, a, b, b', c, c', d denote elements of S , and w denotes a function from $[\text{the carrier of } S, \text{ the carrier of } S]$ into the carrier of G .

Let us consider S, G, w . We say that w is an atlas of S, G if and only if the conditions (Def. 3) are satisfied.

(Def. 3)(i) For all a, x there exists b such that $w(a, b) = x$,

(ii) for all a, b, c such that $w(a, b) = w(a, c)$ holds $b = c$, and

(iii) for all a, b, c holds $w(a, b) + w(b, c) = w(a, c)$.

Let us consider S, G, w, a, x . Let us assume that w is an atlas of S, G . The functor $(a, x).w$ yielding an element of S is defined by:

(Def. 4) $w(a, (a, x).w) = x$.

In the sequel G denotes an add-associative right zeroed right complementable non empty loop structure, x denotes an element of G , and w denotes a function from $[\text{the carrier of } S, \text{ the carrier of } S]$ into the carrier of G .

The following propositions are true:

- (2) $2(0_G) = 0_G$.
- (4)¹ If w is an atlas of S, G , then $w(a, a) = 0_G$.
- (5) If w is an atlas of S, G and $w(a, b) = 0_G$, then $a = b$.
- (6) If w is an atlas of S, G , then $w(a, b) = -w(b, a)$.
- (7) If w is an atlas of S, G and $w(a, b) = w(c, d)$, then $w(b, a) = w(d, c)$.
- (8) If w is an atlas of S, G , then for all b, x there exists a such that $w(a, b) = x$.
- (9) If w is an atlas of S, G and $w(b, a) = w(c, a)$, then $b = c$.
- (10) Let w be a function from $[\text{the carrier of } M, \text{ the carrier of } M]$ into the carrier of G . Suppose w is an atlas of the carrier of M, G and M, G are associated w.r.t. w . Then $p @ q = q @ p$.
- (11) Let w be a function from $[\text{the carrier of } M, \text{ the carrier of } M]$ into the carrier of G . Suppose w is an atlas of the carrier of M, G and M, G are associated w.r.t. w . Then there exists r such that $r @ p = q$.

In the sequel G is an add-associative right zeroed right complementable Abelian non empty loop structure and x is an element of G .

The following propositions are true:

- (13)² Let G be an add-associative Abelian non empty loop structure and x, y, z, t be elements of G . Then $(x + y) + (z + t) = x + z + (y + t)$.
- (14) For every add-associative Abelian non empty loop structure G and for all elements x, y of G holds $2(x + y) = 2x + 2y$.
- (15) $2(-x) = -2x$.
- (16) Let w be a function from $[\text{the carrier of } M, \text{ the carrier of } M]$ into the carrier of G . Suppose w is an atlas of the carrier of M, G and M, G are associated w.r.t. w . Let a, b, c, d be points of M . Then $a @ b = c @ d$ if and only if $w(a, d) = w(c, b)$.

In the sequel w denotes a function from $[\text{the carrier of } M, \text{ the carrier of } M]$ into the carrier of G .

One can prove the following proposition

- (17) If w is an atlas of S, G , then for all a, b, b', c, c' such that $w(a, b) = w(b, c)$ and $w(a, b') = w(b', c')$ holds $w(c, c') = 2w(b, b')$.

We use the following convention: M denotes a midpoint algebra and p, q, r, s denote points of M .

Let us consider M . Observe that $\text{vectgroup}M$ is Abelian, add-associative, right zeroed, and right complementable.

We now state the proposition

- (18)(i) For every set a holds a is an element of $\text{vectgroup}M$ iff a is a vector of M ,
- (ii) $0_{\text{vectgroup}M} = I_M$, and
- (iii) for all elements a, b of $\text{vectgroup}M$ and for all vectors x, y of M such that $a = x$ and $b = y$ holds $a + b = x + y$.

Let I_1 be a non empty loop structure. We say that I_1 is midpoint operator if and only if the conditions (Def. 5) are satisfied.

- (Def. 5)(i) For every element a of I_1 there exists an element x of I_1 such that $2x = a$, and
- (ii) for every element a of I_1 such that $2a = 0_{(I_1)}$ holds $a = 0_{(I_1)}$.

¹ The proposition (3) has been removed.

² The proposition (12) has been removed.

Let us observe that every non empty loop structure which is midpoint operator is also Fanoian.

Let us mention that there exists a non empty loop structure which is strict, midpoint operator, add-associative, right zeroed, right complementable, and Abelian.

In the sequel G denotes a midpoint operator add-associative right zeroed right complementable Abelian non empty loop structure and x, y denote elements of G .

We now state two propositions:

- (19) Let G be a Fanoian add-associative right zeroed right complementable non empty loop structure and x be an element of G . If $x = -x$, then $x = 0_G$.
- (20) Let G be a Fanoian add-associative right zeroed right complementable Abelian non empty loop structure and x, y be elements of G . If $2x = 2y$, then $x = y$.

Let G be a midpoint operator add-associative right zeroed right complementable Abelian non empty loop structure and let x be an element of G . The functor $\frac{1}{2}x$ yielding an element of G is defined as follows:

(Def. 6) $2\frac{1}{2}x = x$.

The following propositions are true:

- (21) $\frac{1}{2}(0_G) = 0_G$ and $\frac{1}{2}(x+y) = \frac{1}{2}x + \frac{1}{2}y$ and if $\frac{1}{2}x = \frac{1}{2}y$, then $x = y$ and $\frac{1}{2}2x = x$.
- (22) Let M be a non empty midpoint algebra structure and w be a function from $[\cdot; \text{the carrier of } M, \text{ the carrier of } M:]$ into the carrier of G . Suppose w is an atlas of the carrier of M, G and M, G are associated w.r.t. w . Let a, b, c, d be points of M . Then $(a @ b) @ (c @ d) = a @ c @ (b @ d)$.
- (23) Let M be a non empty midpoint algebra structure and w be a function from $[\cdot; \text{the carrier of } M, \text{ the carrier of } M:]$ into the carrier of G . Suppose w is an atlas of the carrier of M, G and M, G are associated w.r.t. w . Then M is a midpoint algebra.

Let us consider M . Observe that $\text{vectgroup } M$ is midpoint operator.

Let us consider M, p, q . The functor q^p yielding an element of $\text{vectgroup } M$ is defined as follows:

(Def. 7) $q^p = \overrightarrow{[p, q]}$.

Let us consider M . The functor $\text{vect } M$ yielding a function from $[\cdot; \text{the carrier of } M, \text{ the carrier of } M:]$ into the carrier of $\text{vectgroup } M$ is defined by:

(Def. 8) $(\text{vect } M)(p, q) = \overrightarrow{[p, q]}$.

One can prove the following propositions:

- (24) $\text{vect } M$ is an atlas of the carrier of $M, \text{vectgroup } M$.
- (25) $\overrightarrow{[p, q]} = \overrightarrow{[r, s]}$ iff $p @ s = q @ r$.
- (26) $p @ q = r$ iff $\overrightarrow{[p, r]} = \overrightarrow{[r, q]}$.
- (27) $M, \text{vectgroup } M$ are associated w.r.t. $\text{vect } M$.

In the sequel w is a function from $[\cdot; S, S:]$ into the carrier of G .

Let us consider S, G, w . Let us assume that w is an atlas of S, G . The functor $@ w$ yielding a binary operation on S is defined as follows:

(Def. 9) $w(a, (@ w)(a, b)) = w((@ w)(a, b), b)$.

We now state the proposition

- (28) If w is an atlas of S, G , then for all a, b, c holds $(@ w)(a, b) = c$ iff $w(a, c) = w(c, b)$.

Let D be a non empty set and let M be a binary operation on D . Observe that $\langle D, M \rangle$ is non empty.

Let us consider S, G, w . The functor $\text{Atlas } w$ yields a function from $[\text{the carrier of } \langle S, @w \rangle, \text{ the carrier of } \langle S, @w \rangle]$ into the carrier of G and is defined by:

(Def. 10) $\text{Atlas } w = w$.

We now state the proposition

(32)³ If w is an atlas of S, G , then $\langle S, @w \rangle, G$ are associated w.r.t. $\text{Atlas } w$.

Let us consider S, G, w . Let us assume that w is an atlas of S, G . The functor $\text{MidSp}(w)$ yields a strict midpoint algebra and is defined by:

(Def. 11) $\text{MidSp}(w) = \langle S, @w \rangle$.

We adopt the following rules: M is a non empty midpoint algebra structure, w is a function from $[\text{the carrier of } M, \text{ the carrier of } M]$ into the carrier of G , and a, b, b_1, b_2, c are points of M .

Next we state the proposition

(33) M is a midpoint algebra if and only if there exists G and there exists w such that w is an atlas of the carrier of M, G and M, G are associated w.r.t. w .

Let M be a non empty midpoint algebra structure. We introduce atlas structures over M which are systems

$\langle \text{an algebra, a function} \rangle$,

where the algebra is a non empty loop structure and the function is a function from $[\text{the carrier of } M, \text{ the carrier of } M]$ into the carrier of the algebra.

Let M be a non empty midpoint algebra structure and let I_1 be an atlas structure over M . We say that I_1 is atlas-like if and only if the conditions (Def. 12) are satisfied.

(Def. 12)(i) The algebra of I_1 is midpoint operator, add-associative, right zeroed, right complementable, and Abelian,

(ii) M , the algebra of I_1 are associated w.r.t. the function of I_1 , and

(iii) the function of I_1 is an atlas of the carrier of M , the algebra of I_1 .

Let M be a midpoint algebra. One can check that there exists an atlas structure over M which is atlas-like.

Let M be a non empty midpoint algebra. An atlas of M is an atlas-like atlas structure over M .

Let M be a non empty midpoint algebra structure and let W be an atlas structure over M . A vector of W is an element of the algebra of W .

Let M be a midpoint algebra, let W be an atlas structure over M , and let a, b be points of M . The functor $W(a, b)$ yields an element of the algebra of W and is defined by:

(Def. 13) $W(a, b) = (\text{the function of } W)(a, b)$.

Let M be a midpoint algebra, let W be an atlas structure over M , let a be a point of M , and let x be a vector of W . The functor $(a, x).W$ yields a point of M and is defined as follows:

(Def. 14) $(a, x).W = (a, x).\text{the function of } W$.

Let M be a midpoint algebra and let W be an atlas of M . The functor 0_W yields a vector of W and is defined as follows:

(Def. 15) $0_W = 0_{\text{the algebra of } W}$.

We now state two propositions:

(34) Suppose w is an atlas of the carrier of M, G and M, G are associated w.r.t. w . Then $a @ c = b_1 @ b_2$ if and only if $w(a, c) = w(a, b_1) + w(a, b_2)$.

³ The propositions (29)–(31) have been removed.

- (35) Suppose w is an atlas of the carrier of M , G and M , G are associated w.r.t. w . Then $a^{\textcircled{}} c = b$ if and only if $w(a, c) = 2w(a, b)$.

For simplicity, we adopt the following rules: M denotes a midpoint algebra, W denotes an atlas of M , a, b, b_1, b_2, c, d denote points of M , and x denotes a vector of W .

We now state several propositions:

- (36) $a^{\textcircled{}} c = b_1^{\textcircled{}} b_2$ iff $W(a, c) = W(a, b_1) + W(a, b_2)$.
- (37) $a^{\textcircled{}} c = b$ iff $W(a, c) = 2W(a, b)$.
- (38)(i) For all a, x there exists b such that $W(a, b) = x$,
(ii) for all a, b, c such that $W(a, b) = W(a, c)$ holds $b = c$, and
(iii) for all a, b, c holds $W(a, b) + W(b, c) = W(a, c)$.
- (39) $W(a, a) = 0_W$ and if $W(a, b) = 0_W$, then $a = b$ and $W(a, b) = -W(b, a)$ and if $W(a, b) = W(c, d)$, then $W(b, a) = W(d, c)$ and for all b, x there exists a such that $W(a, b) = x$ and if $W(b, a) = W(c, a)$, then $b = c$ and $a^{\textcircled{}} b = c$ iff $W(a, c) = W(c, b)$ and $a^{\textcircled{}} b = c^{\textcircled{}} d$ iff $W(a, d) = W(c, b)$ and $W(a, b) = x$ iff $(a, x).W = b$.
- (40) $(a, 0_W).W = a$.

REFERENCES

- [1] Czesław Byliński. Binary operations. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/binop_1.html.
- [2] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/funct_2.html.
- [3] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/zfmisc_1.html.
- [4] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/vectsp_1.html.
- [5] Michał Muzalewski. Midpoint algebras. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/midsp_1.html.
- [6] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/rivect_1.html.
- [7] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/subset_1.html.

Received June 21, 1991

Published January 2, 2004