

# Free Order Sorted Universal Algebra<sup>1</sup>

Josef Urban  
Charles University  
Praha

**Summary.** Free Order Sorted Universal Algebra — the general construction for any locally directed signatures.

MML Identifier: OSAFREE.

WWW: <http://mizar.org/JFM/Vol14/osafree.html>

The articles [22], [13], [28], [33], [34], [10], [23], [12], [11], [7], [14], [35], [4], [19], [2], [21], [27], [15], [5], [3], [6], [1], [8], [26], [24], [18], [25], [9], [16], [17], [30], [32], [29], [31], and [20] provide the notation and terminology for this paper.

## 1. PRELIMINARIES

In this paper  $S$  is an order sorted signature.

Let  $S$  be an order sorted signature and let  $U_0$  be an order sorted algebra of  $S$ . A subset of  $U_0$  is called an order sorted generator set of  $U_0$  if:

(Def. 1) For every OSSubset  $O$  of  $U_0$  such that  $O = \text{OSCl}$  it holds the sorts of  $\text{OSGen } O =$  the sorts of  $U_0$ .

One can prove the following proposition

(1) Let  $S$  be an order sorted signature,  $U_0$  be a strict non-empty order sorted algebra of  $S$ , and  $A$  be a subset of  $U_0$ . Then  $A$  is an order sorted generator set of  $U_0$  if and only if for every OSSubset  $O$  of  $U_0$  such that  $O = \text{OSCl } A$  holds  $\text{OSGen } O = U_0$ .

Let us consider  $S$ , let  $U_0$  be a monotone order sorted algebra of  $S$ , and let  $I_1$  be an order sorted generator set of  $U_0$ . We say that  $I_1$  is osfree if and only if the condition (Def. 2) is satisfied.

(Def. 2) Let  $U_1$  be a monotone non-empty order sorted algebra of  $S$  and  $f$  be a many sorted function from  $I_1$  into the sorts of  $U_1$ . Then there exists a many sorted function  $h$  from  $U_0$  into  $U_1$  such that  $h$  is a homomorphism of  $U_0$  into  $U_1$  and order-sorted and  $h \upharpoonright I_1 = f$ .

Let  $S$  be an order sorted signature and let  $I_1$  be a monotone order sorted algebra of  $S$ . We say that  $I_1$  is osfree if and only if:

(Def. 3) There exists an order sorted generator set of  $I_1$  which is osfree.

---

<sup>1</sup>This work was done during author's research visit in Białystok, funded by the CALCULEMUS grant HPRN-CT-2000-00102.

## 2. CONSTRUCTION OF FREE ORDER SORTED ALGEBRAS FOR GIVEN SIGNATURE

Let  $S$  be an order sorted signature and let  $X$  be a many sorted set indexed by  $S$ . The functor  $\text{OSRELX}$  yields a relation between  $[\text{the operation symbols of } S, \{\text{the carrier of } S\}] \cup \bigcup \text{coprod}(X)$  and  $([\text{the operation symbols of } S, \{\text{the carrier of } S\}] \cup \bigcup \text{coprod}(X))^*$  and is defined by the condition (Def. 4).

(Def. 4) Let  $a$  be an element of  $[\text{the operation symbols of } S, \{\text{the carrier of } S\}] \cup \bigcup \text{coprod}(X)$  and  $b$  be an element of  $([\text{the operation symbols of } S, \{\text{the carrier of } S\}] \cup \bigcup \text{coprod}(X))^*$ . Then  $\langle a, b \rangle \in \text{OSRELX}$  if and only if the following conditions are satisfied:

- (i)  $a \in [\text{the operation symbols of } S, \{\text{the carrier of } S\}]$ , and
- (ii) for every operation symbol  $o$  of  $S$  such that  $\langle o, \text{the carrier of } S \rangle = a$  holds  $\text{len } b = \text{len Arity}(o)$  and for every set  $x$  such that  $x \in \text{dom } b$  holds if  $b(x) \in [\text{the operation symbols of } S, \{\text{the carrier of } S\}]$ , then for every operation symbol  $o_1$  of  $S$  such that  $\langle o_1, \text{the carrier of } S \rangle = b(x)$  holds the result sort of  $o_1 \leq \text{Arity}(o)_x$  and if  $b(x) \in \bigcup \text{coprod}(X)$ , then there exists an element  $i$  of  $S$  such that  $i \leq \text{Arity}(o)_x$  and  $b(x) \in \text{coprod}(i, X)$ .

In the sequel  $S$  denotes an order sorted signature,  $X$  denotes a many sorted set indexed by  $S$ ,  $o$  denotes an operation symbol of  $S$ , and  $b$  denotes an element of  $([\text{the operation symbols of } S, \{\text{the carrier of } S\}] \cup \bigcup \text{coprod}(X))^*$ .

We now state the proposition

(2)  $\langle \langle o, \text{the carrier of } S \rangle, b \rangle \in \text{OSRELX}$  if and only if the following conditions are satisfied:

- (i)  $\text{len } b = \text{len Arity}(o)$ , and
- (ii) for every set  $x$  such that  $x \in \text{dom } b$  holds if  $b(x) \in [\text{the operation symbols of } S, \{\text{the carrier of } S\}]$ , then for every operation symbol  $o_1$  of  $S$  such that  $\langle o_1, \text{the carrier of } S \rangle = b(x)$  holds the result sort of  $o_1 \leq \text{Arity}(o)_x$  and if  $b(x) \in \bigcup \text{coprod}(X)$ , then there exists an element  $i$  of  $S$  such that  $i \leq \text{Arity}(o)_x$  and  $b(x) \in \text{coprod}(i, X)$ .

Let  $S$  be an order sorted signature and let  $X$  be a many sorted set indexed by  $S$ . The functor  $\text{DTConOSAX}$  yields a tree construction structure and is defined by:

(Def. 5)  $\text{DTConOSAX} = \langle [\text{the operation symbols of } S, \{\text{the carrier of } S\}] \cup \bigcup \text{coprod}(X), \text{OSRELX} \rangle$ .

Let  $S$  be an order sorted signature and let  $X$  be a many sorted set indexed by  $S$ . Note that  $\text{DTConOSAX}$  is strict and non empty.

The following proposition is true

- (3) Let  $S$  be an order sorted signature and  $X$  be a non-empty many sorted set indexed by  $S$ . Then the nonterminals of  $\text{DTConOSAX} = [\text{the operation symbols of } S, \{\text{the carrier of } S\}]$  and the terminals of  $\text{DTConOSAX} = \bigcup \text{coprod}(X)$ .

Let  $S$  be an order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . Note that  $\text{DTConOSAX}$  has terminals, nonterminals, and useful nonterminals.

We now state the proposition

- (4) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $t$  be a set. Then  $t \in$  the terminals of  $\text{DTConOSAX}$  if and only if there exists an element  $s$  of  $S$  and there exists a set  $x$  such that  $x \in X(s)$  and  $t = \langle x, s \rangle$ .

Let  $S$  be an order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $o$  be an operation symbol of  $S$ . The functor  $\text{OSSym}(o, X)$  yields a symbol of  $\text{DTConOSAX}$  and is defined as follows:

(Def. 6)  $\text{OSSym}(o, X) = \langle o, \text{the carrier of } S \rangle$ .

Let  $S$  be an order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $s$  be an element of  $S$ . The functor  $\text{ParsedTerms}(X, s)$  yields a subset of  $\text{TS}(\text{DTConOSAX})$  and is defined by the condition (Def. 7).

(Def. 7)  $\text{ParsedTerms}(X, s) = \{a; a \text{ ranges over elements of } \text{TS}(\text{DTConOSAX}): \bigvee_{s_1: \text{element of } S} \bigvee_{x: \text{set } (s_1 \leq s \wedge x \in X(s_1))} a = \text{the root tree of } \langle x, s_1 \rangle\} \vee \bigvee_{o: \text{operation symbol of } S} (\langle o, \text{the carrier of } S \rangle = a(\emptyset) \wedge \text{the result sort of } o \leq s)$ .

Let  $S$  be an order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $s$  be an element of  $S$ . Observe that  $\text{ParsedTerms}(X, s)$  is non empty.

Let  $S$  be an order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $\text{ParsedTerms}X$  yields an order sorted set of  $S$  and is defined by:

(Def. 8) For every element  $s$  of  $S$  holds  $(\text{ParsedTerms}X)(s) = \text{ParsedTerms}(X, s)$ .

Let  $S$  be an order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . Observe that  $\text{ParsedTerms}X$  is non-empty.

Next we state four propositions:

- (5) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $o$  be an operation symbol of  $S$ , and  $x$  be a set. Suppose  $x \in ((\text{ParsedTerms}X)^\# \cdot \text{the arity of } S)(o)$ . Then  $x$  is a finite sequence of elements of  $\text{TS}(\text{DTConOSAX})$ .
- (6) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $o$  be an operation symbol of  $S$ , and  $p$  be a finite sequence of elements of  $\text{TS}(\text{DTConOSAX})$ . Then  $p \in ((\text{ParsedTerms}X)^\# \cdot \text{the arity of } S)(o)$  if and only if  $\text{dom } p = \text{dom Arity}(o)$  and for every natural number  $n$  such that  $n \in \text{dom } p$  holds  $p(n) \in \text{ParsedTerms}(X, \text{Arity}(o)_n)$ .
- (7) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $o$  be an operation symbol of  $S$ , and  $p$  be a finite sequence of elements of  $\text{TS}(\text{DTConOSAX})$ . Then  $\text{OSSym}(o, X) \Rightarrow \text{the roots of } p$  if and only if  $p \in ((\text{ParsedTerms}X)^\# \cdot \text{the arity of } S)(o)$ .
- (8) For every order sorted signature  $S$  and for every non-empty many sorted set  $X$  indexed by  $S$  holds  $\bigcup \text{rng } \text{ParsedTerms}X = \text{TS}(\text{DTConOSAX})$ .

Let  $S$  be an order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $o$  be an operation symbol of  $S$ . The functor  $\text{PTDenOp}(o, X)$  yields a function from  $((\text{ParsedTerms}X)^\# \cdot \text{the arity of } S)(o)$  into  $(\text{ParsedTerms}X \cdot \text{the result sort of } S)(o)$  and is defined by:

(Def. 9) For every finite sequence  $p$  of elements of  $\text{TS}(\text{DTConOSAX})$  such that  $\text{OSSym}(o, X) \Rightarrow \text{the roots of } p$  holds  $(\text{PTDenOp}(o, X))(p) = \text{OSSym}(o, X)\text{-tree}(p)$ .

Let  $S$  be an order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $\text{PTOper}X$  yields a many sorted function from  $(\text{ParsedTerms}X)^\# \cdot \text{the arity of } S$  into  $\text{ParsedTerms}X \cdot \text{the result sort of } S$  and is defined as follows:

(Def. 10) For every operation symbol  $o$  of  $S$  holds  $(\text{PTOper}X)(o) = \text{PTDenOp}(o, X)$ .

Let  $S$  be an order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $\text{ParsedTermsOSAX}$  yields an order sorted algebra of  $S$  and is defined by:

(Def. 11)  $\text{ParsedTermsOSAX} = \langle \text{ParsedTerms}X, \text{PTOper}X \rangle$ .

Let  $S$  be an order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . Note that  $\text{ParsedTermsOSAX}$  is strict and non-empty.

Let  $S$  be an order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $o$  be an operation symbol of  $S$ . Then  $\text{OSSym}(o, X)$  is a nonterminal of  $\text{DTConOSAX}$ .

One can prove the following propositions:

- (9) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $s$  be an element of  $S$ . Then (the sorts of  $\text{ParsedTermsOSAX}$ )( $s$ ) =  $\{a; a \text{ ranges over elements of } \text{TS}(\text{DTConOSAX}): \bigvee_{s_1: \text{element of } S} \bigvee_{x: \text{set}} (s_1 \leq s \wedge x \in X(s_1) \wedge a = \text{the root tree of } \langle x, s_1 \rangle) \vee \bigvee_{o: \text{operation symbol of } S} (\langle o, \text{the carrier of } S \rangle = a(\emptyset) \wedge \text{the result sort of } o \leq s)\}$ .
- (10) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $s, s_1$  be elements of  $S$ , and  $x$  be a set. Suppose  $x \in X(s)$ . Then
- (i) the root tree of  $\langle x, s \rangle$  is an element of  $\text{TS}(\text{DTConOSAX})$ ,
  - (ii) for every set  $z$  holds  $\langle z, \text{the carrier of } S \rangle \neq (\text{the root tree of } \langle x, s \rangle)(\emptyset)$ , and
  - (iii) the root tree of  $\langle x, s \rangle \in (\text{the sorts of } \text{ParsedTermsOSAX})(s_1)$  iff  $s \leq s_1$ .
- (11) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ , and  $o$  be an operation symbol of  $S$ . Suppose  $t(\emptyset) = \langle o, \text{the carrier of } S \rangle$ . Then
- (i) there exists a subtree sequence  $p$  joinable by  $\text{OSSym}(o, X)$  such that  $t = \text{OSSym}(o, X)\text{-tree}(p)$  and  $\text{OSSym}(o, X) \Rightarrow$  the roots of  $p$  and  $p \in \text{Args}(o, \text{ParsedTermsOSAX})$  and  $t = (\text{Den}(o, \text{ParsedTermsOSAX}))(p)$ ,
  - (ii) for every element  $s_2$  of  $S$  and for every set  $x$  holds  $t \neq$  the root tree of  $\langle x, s_2 \rangle$ , and
  - (iii) for every element  $s_1$  of  $S$  holds  $t \in (\text{the sorts of } \text{ParsedTermsOSAX})(s_1)$  iff the result sort of  $o \leq s_1$ .
- (12) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $n_1$  be a symbol of  $\text{DTConOSAX}$ , and  $t_1$  be a finite sequence of elements of  $\text{TS}(\text{DTConOSAX})$ . Suppose  $n_1 \Rightarrow$  the roots of  $t_1$ . Then
- (i)  $n_1 \in$  the nonterminals of  $\text{DTConOSAX}$ ,
  - (ii)  $n_1\text{-tree}(t_1) \in \text{TS}(\text{DTConOSAX})$ , and
  - (iii) there exists an operation symbol  $o$  of  $S$  such that  $n_1 = \langle o, \text{the carrier of } S \rangle$  and  $t_1 \in \text{Args}(o, \text{ParsedTermsOSAX})$  and  $n_1\text{-tree}(t_1) = (\text{Den}(o, \text{ParsedTermsOSAX}))(t_1)$  and for every element  $s_1$  of  $S$  holds  $n_1\text{-tree}(t_1) \in (\text{the sorts of } \text{ParsedTermsOSAX})(s_1)$  iff the result sort of  $o \leq s_1$ .
- (13) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $o$  be an operation symbol of  $S$ , and  $x$  be a finite sequence. Then  $x \in \text{Args}(o, \text{ParsedTermsOSAX})$  if and only if the following conditions are satisfied:
- (i)  $x$  is a finite sequence of elements of  $\text{TS}(\text{DTConOSAX})$ , and
  - (ii)  $\text{OSSym}(o, X) \Rightarrow$  the roots of  $x$ .
- (14) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ . Then there exists a sort symbol  $s$  of  $S$  such that  $t \in (\text{the sorts of } \text{ParsedTermsOSAX})(s)$  and for every element  $s_1$  of  $S$  such that  $t \in (\text{the sorts of } \text{ParsedTermsOSAX})(s_1)$  holds  $s \leq s_1$ .

Let  $S$  be an order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ . The functor  $\text{LeastSort}t$  yields a sort symbol of  $S$  and is defined by:

- (Def. 12)  $t \in (\text{the sorts of } \text{ParsedTermsOSAX})(\text{LeastSort}t)$  and for every element  $s_1$  of  $S$  such that  $t \in (\text{the sorts of } \text{ParsedTermsOSAX})(s_1)$  holds  $\text{LeastSort}t \leq s_1$ .

Let  $S$  be a non empty non void many sorted signature and let  $A$  be a non-empty algebra over  $S$ . An element of  $A$  is an element of  $\bigcup$ (the sorts of  $A$ ).

One can prove the following propositions:

- (15) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $x$  be a set. Then  $x$  is an element of  $\text{ParsedTermsOSAX}$  if and only if  $x$  is an element of  $\text{TS}(\text{DTConOSAX})$ .

- (16) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $s$  be an element of  $S$ , and  $x$  be a set. If  $x \in (\text{the sorts of } \text{ParsedTermsOSAX})(s)$ , then  $x$  is an element of  $\text{TS}(\text{DTConOSAX})$ .
- (17) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $s$  be an element of  $S$ , and  $x$  be a set. Suppose  $x \in X(s)$ . Let  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ . If  $t = \text{the root tree of } \langle x, s \rangle$ , then  $\text{LeastSort } t = s$ .
- (18) Let  $S$  be an order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $o$  be an operation symbol of  $S$ ,  $x$  be an element of  $\text{Args}(o, \text{ParsedTermsOSAX})$ , and  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ . If  $t = (\text{Den}(o, \text{ParsedTermsOSAX}))(x)$ , then  $\text{LeastSort } t = \text{the result sort of } o$ .

Let  $S$  be an order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $o_2$  be an operation symbol of  $S$ . Note that  $\text{Args}(o_2, \text{ParsedTermsOSAX})$  is non empty.

Let  $S$  be a locally directed order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $x$  be a finite sequence of elements of  $\text{TS}(\text{DTConOSAX})$ . The functor  $\text{LeastSorts } x$  yielding an element of  $(\text{the carrier of } S)^*$  is defined by:

(Def. 14)<sup>1</sup>  $\text{dom } \text{LeastSorts } x = \text{dom } x$  and for every natural number  $y$  such that  $y \in \text{dom } x$  there exists an element  $t$  of  $\text{TS}(\text{DTConOSAX})$  such that  $t = x(y)$  and  $(\text{LeastSorts } x)(y) = \text{LeastSort } t$ .

One can prove the following proposition

- (19) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $o$  be an operation symbol of  $S$ , and  $x$  be a finite sequence of elements of  $\text{TS}(\text{DTConOSAX})$ . Then  $\text{LeastSorts } x \leq \text{Arity}(o)$  if and only if  $x \in \text{Args}(o, \text{ParsedTermsOSAX})$ .

One can verify that there exists a monotone order sorted signature which is locally directed and regular.

Let  $S$  be a locally directed regular monotone order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , let  $o$  be an operation symbol of  $S$ , and let  $x$  be a finite sequence of elements of  $\text{TS}(\text{DTConOSAX})$ . Let us assume that  $\text{OSSym}(\text{LBound}(o, \text{LeastSorts } x), X) \Rightarrow \text{the roots of } x$ . The functor  $\pi_x o$  yielding an element of  $\text{TS}(\text{DTConOSAX})$  is defined as follows:

(Def. 15)  $\pi_x o = \text{OSSym}(\text{LBound}(o, \text{LeastSorts } x), X)\text{-tree}(x)$ .

Let  $S$  be a locally directed order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $t$  be a symbol of  $\text{DTConOSAX}$ . Let us assume that there exists a finite sequence  $p$  such that  $t \Rightarrow p$ . The functor  ${}^@ (X, t)$  yielding an operation symbol of  $S$  is defined by:

(Def. 16)  $\langle {}^@ (X, t), \text{the carrier of } S \rangle = t$ .

Let  $S$  be an order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $t$  be a symbol of  $\text{DTConOSAX}$ . Let us assume that  $t \in \text{the terminals of } \text{DTConOSAX}$ . The functor  $\prod t$  yielding an element of  $\text{TS}(\text{DTConOSAX})$  is defined by:

(Def. 17)  $\prod t = \text{the root tree of } t$ .

Let  $S$  be a locally directed order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $\text{LCongruence } X$  yields a monotone order sorted congruence of  $\text{ParsedTermsOSAX}$  and is defined by:

(Def. 18) For every monotone order sorted congruence  $R$  of  $\text{ParsedTermsOSAX}$  holds  $\text{LCongruence } X \subseteq R$ .

Let  $S$  be a locally directed order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $\text{FreeOSAX}$  yielding a strict non-empty monotone order sorted algebra of  $S$  is defined by:

<sup>1</sup> The definition (Def. 13) has been removed.

(Def. 19)  $\text{FreeOSAX} = \text{QuotOSAlg}(\text{ParsedTermsOSAX}, \text{LCongruenceX})$ .

Let  $S$  be an order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $t$  be a symbol of  $\text{DTConOSAX}$ . The functor  ${}^@t$  yielding a subset of  $[\text{TS}(\text{DTConOSAX})]$ , the carrier of  $S$  is defined by the condition (Def. 20).

(Def. 20)  ${}^@t = \{\langle \text{the root tree of } t, s_1 \rangle; s_1 \text{ ranges over elements of } S: \bigvee_{s: \text{element of } S} \bigvee_{x: \text{set}} (x \in X(s) \wedge t = \langle x, s \rangle \wedge s \leq s_1)\}$ .

Let  $S$  be an order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , let  $n_1$  be a symbol of  $\text{DTConOSAX}$ , and let  $x$  be a finite sequence of elements of  $[\text{TS}(\text{DTConOSAX})]$ , the carrier of  $S$ . The functor  ${}^@(n_1, x)$  yielding a subset of  $[\text{TS}(\text{DTConOSAX})]$ , the carrier of  $S$  is defined by the condition (Def. 21).

(Def. 21)  ${}^@(n_1, x) = \{\langle (\text{Den}(o_2, \text{ParsedTermsOSAX}))(x_2), s_3 \rangle; o_2 \text{ ranges over operation symbols of } S, x_2 \text{ ranges over elements of } \text{Args}(o_2, \text{ParsedTermsOSAX}), s_3 \text{ ranges over elements of } S: \bigvee_{o_1: \text{operation symbol of } S} (n_1 = \langle o_1, \text{the carrier of } S \rangle \wedge o_1 \cong o_2 \wedge \text{len Arity}(o_1) = \text{len Arity}(o_2) \wedge \text{the result sort of } o_1 \leq s_3 \wedge \text{the result sort of } o_2 \leq s_3) \wedge \bigvee_{w_3: \text{element of (the carrier of } S)^*} (\text{dom } w_3 = \text{dom } x \wedge \bigwedge_{y: \text{natural number}} (y \in \text{dom } x \Rightarrow \langle x_2(y), (w_3)_y \rangle \in x(y))\}$ .

Let  $S$  be a locally directed order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $\text{PTClassesX}$  yielding a function from  $\text{TS}(\text{DTConOSAX})$  into  $[\text{TS}(\text{DTConOSAX})]$ , the carrier of  $S$  is defined by the conditions (Def. 22).

(Def. 22)(i) For every symbol  $t$  of  $\text{DTConOSAX}$  such that  $t \in$  the terminals of  $\text{DTConOSAX}$  holds  $(\text{PTClassesX})(\text{the root tree of } t) = {}^@t$ , and  
(ii) for every symbol  $n_1$  of  $\text{DTConOSAX}$  and for every finite sequence  $t_1$  of elements of  $\text{TS}(\text{DTConOSAX})$  such that  $n_1 \Rightarrow$  the roots of  $t_1$  holds  $(\text{PTClassesX})(n_1\text{-tree}(t_1)) = {}^@(n_1, \text{PTClassesX} \cdot t_1)$ .

Next we state four propositions:

(20) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ . Then

- (i) for every element  $s$  of  $S$  holds  $t \in$  (the sorts of  $\text{ParsedTermsOSAX})(s)$  iff  $\langle t, s \rangle \in (\text{PTClassesX})(t)$ , and
- (ii) for every element  $s$  of  $S$  and for every element  $y$  of  $\text{TS}(\text{DTConOSAX})$  such that  $\langle y, s \rangle \in (\text{PTClassesX})(t)$  holds  $\langle t, s \rangle \in (\text{PTClassesX})(y)$ .

(21) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ , and  $s$  be an element of  $S$ . If there exists an element  $y$  of  $\text{TS}(\text{DTConOSAX})$  such that  $\langle y, s \rangle \in (\text{PTClassesX})(t)$ , then  $\langle t, s \rangle \in (\text{PTClassesX})(t)$ .

(22) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $x, y$  be elements of  $\text{TS}(\text{DTConOSAX})$ , and  $s_1, s_2$  be elements of  $S$ . Suppose  $s_1 \leq s_2$  and  $x \in$  (the sorts of  $\text{ParsedTermsOSAX})(s_1)$  and  $y \in$  (the sorts of  $\text{ParsedTermsOSAX})(s_1)$ . Then  $\langle y, s_1 \rangle \in (\text{PTClassesX})(x)$  if and only if  $\langle y, s_2 \rangle \in (\text{PTClassesX})(x)$ .

(23) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $x, y, z$  be elements of  $\text{TS}(\text{DTConOSAX})$ , and  $s$  be an element of  $S$ . If  $\langle y, s \rangle \in (\text{PTClassesX})(x)$  and  $\langle z, s \rangle \in (\text{PTClassesX})(y)$ , then  $\langle x, s \rangle \in (\text{PTClassesX})(z)$ .

Let  $S$  be a locally directed order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $\text{PTCongruenceX}$  yields an equivalence order sorted relation of  $\text{ParsedTermsOSAX}$  and is defined by the condition (Def. 23).

(Def. 23) Let  $i$  be a set. Suppose  $i \in$  the carrier of  $S$ . Then  $(\text{PTCongruenceX})(i) = \{\langle x, y \rangle; x \text{ ranges over elements of } \text{TS}(\text{DTConOSAX}), y \text{ ranges over elements of } \text{TS}(\text{DTConOSAX}): \langle x, i \rangle \in (\text{PTClassesX})(y)\}$ .

The following propositions are true:

- (24) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $x, y, s$  be sets. If  $\langle x, s \rangle \in (\text{PTClasses}X)(y)$ , then  $x \in \text{TS}(\text{DTConOSAX})$  and  $y \in \text{TS}(\text{DTConOSAX})$  and  $s \in$  the carrier of  $S$ .
- (25) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $C$  be a component of  $S$ , and  $x, y$  be sets. Then  $\langle x, y \rangle \in \text{CompClass}(\text{PTCongruence}X, C)$  if and only if there exists an element  $s_1$  of  $S$  such that  $s_1 \in C$  and  $\langle x, s_1 \rangle \in (\text{PTClasses}X)(y)$ .
- (26) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $s$  be an element of  $S$ , and  $x$  be an element of (the sorts of  $\text{ParsedTermsOSAX})(s)$ . Then  $\text{OSClass}(\text{PTCongruence}X, x) = \pi_1((\text{PTClasses}X)(x))$ .
- (27) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $R$  be a many sorted relation indexed by  $\text{ParsedTermsOSAX}$ . Then  $R = \text{PTCongruence}X$  if and only if the following conditions are satisfied:
- (i) for all elements  $s_1, s_2$  of  $S$  and for every set  $x$  such that  $x \in X(s_1)$  holds if  $s_1 \leq s_2$ , then (the root tree of  $\langle x, s_1 \rangle$ , the root tree of  $\langle x, s_1 \rangle \in R(s_2)$  and for every set  $y$  such that (the root tree of  $\langle x, s_1 \rangle, y \in R(s_2)$  or  $\langle y, \text{the root tree of } \langle x, s_1 \rangle \rangle \in R(s_2)$  holds  $s_1 \leq s_2$  and  $y =$  the root tree of  $\langle x, s_1 \rangle$ , and
  - (ii) for all operation symbols  $o_1, o_2$  of  $S$  and for every element  $x_1$  of  $\text{Args}(o_1, \text{ParsedTermsOSAX})$  and for every element  $x_2$  of  $\text{Args}(o_2, \text{ParsedTermsOSAX})$  and for every element  $s_3$  of  $S$  holds  $\langle (\text{Den}(o_1, \text{ParsedTermsOSAX}))(x_1), (\text{Den}(o_2, \text{ParsedTermsOSAX}))(x_2) \rangle \in R(s_3)$  iff  $o_1 \cong o_2$  and  $\text{lenArity}(o_1) = \text{lenArity}(o_2)$  and the result sort of  $o_1 \leq s_3$  and the result sort of  $o_2 \leq s_3$  and there exists an element  $w_3$  of (the carrier of  $S$ )\* such that  $\text{dom } w_3 = \text{dom } x_1$  and for every natural number  $y$  such that  $y \in \text{dom } w_3$  holds  $\langle x_1(y), x_2(y) \rangle \in R((w_3)_y)$ .
- (28) Let  $S$  be a locally directed order sorted signature and  $X$  be a non-empty many sorted set indexed by  $S$ . Then  $\text{PTCongruence}X$  is monotone.

Let  $S$  be a locally directed order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . Observe that  $\text{PTCongruence}X$  is monotone.

Let  $S$  be a locally directed order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $s$  be an element of  $S$ . The functor  $\text{PTVars}(s, X)$  yields a subset of (the sorts of  $\text{ParsedTermsOSAX})(s)$  and is defined as follows:

- (Def. 24) For every set  $x$  holds  $x \in \text{PTVars}(s, X)$  iff there exists a set  $a$  such that  $a \in X(s)$  and  $x =$  the root tree of  $\langle a, s \rangle$ .

Let  $S$  be a locally directed order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $s$  be an element of  $S$ . Note that  $\text{PTVars}(s, X)$  is non empty.

Next we state the proposition

- (29) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $s$  be an element of  $S$ . Then  $\text{PTVars}(s, X) = \{\text{the root tree of } t; t \text{ ranges over symbols of } \text{DTConOSAX} : t \in \text{the terminals of } \text{DTConOSAX} \wedge t_2 = s\}$ .

Let  $S$  be a locally directed order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $\text{PTVars}X$  yielding a subset of  $\text{ParsedTermsOSAX}$  is defined by:

- (Def. 25) For every element  $s$  of  $S$  holds  $(\text{PTVars}X)(s) = \text{PTVars}(s, X)$ .

One can prove the following proposition

- (30) Let  $S$  be a locally directed order sorted signature and  $X$  be a non-empty many sorted set indexed by  $S$ . Then  $\text{PTVars}X$  is non-empty.

Let  $S$  be a locally directed order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $s$  be an element of  $S$ . The functor  $\text{OSFreeGen}(s, X)$  yielding a subset of (the sorts of  $\text{FreeOSAX}$ )( $s$ ) is defined as follows:

- (Def. 26) For every set  $x$  holds  $x \in \text{OSFreeGen}(s, X)$  iff there exists a set  $a$  such that  $a \in X(s)$  and  $x = (\text{OSNatHom}(\text{ParsedTermsOSAX}, \text{LCongruenceX})) (s)$  (the root tree of  $\langle a, s \rangle$ ).

Let  $S$  be a locally directed order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $s$  be an element of  $S$ . One can verify that  $\text{OSFreeGen}(s, X)$  is non empty.

Next we state the proposition

- (31) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $s$  be an element of  $S$ . Then  $\text{OSFreeGen}(s, X) = \{(\text{OSNatHom}(\text{ParsedTermsOSAX}, \text{LCongruenceX})) (s)$  (the root tree of  $t$ );  $t$  ranges over symbols of  $\text{DTConOSAX} : t \in \text{the terminals of DTConOSAX} \wedge t_2 = s\}$ .

Let  $S$  be a locally directed order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $\text{OSFreeGenX}$  yields an order sorted generator set of  $\text{FreeOSAX}$  and is defined by:

- (Def. 27) For every element  $s$  of  $S$  holds  $(\text{OSFreeGenX})(s) = \text{OSFreeGen}(s, X)$ .

The following proposition is true

- (32) Let  $S$  be a locally directed order sorted signature and  $X$  be a non-empty many sorted set indexed by  $S$ . Then  $\text{OSFreeGenX}$  is non-empty.

Let  $S$  be a locally directed order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . One can check that  $\text{OSFreeGenX}$  is non-empty.

Let  $S$  be a locally directed order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , let  $R$  be an order sorted congruence of  $\text{ParsedTermsOSAX}$ , and let  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ . The functor  $\text{OSClass}(R, t)$  yields an element of  $\text{OSClass}(R, \text{LeastSort } t)$  and is defined as follows:

- (Def. 28) For every element  $s$  of  $S$  and for every element  $x$  of (the sorts of  $\text{ParsedTermsOSAX}$ )( $s$ ) such that  $t = x$  holds  $\text{OSClass}(R, t) = \text{OSClass}(R, x)$ .

Next we state several propositions:

- (33) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $R$  be an order sorted congruence of  $\text{ParsedTermsOSAX}$ , and  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ . Then  $t \in \text{OSClass}(R, t)$ .

- (34) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $s$  be an element of  $S$ ,  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ , and  $x, x_1$  be sets. Suppose  $x \in X(s)$  and  $t$  = the root tree of  $\langle x, s \rangle$ . Then  $x_1 \in \text{OSClass}(\text{PTCongruenceX}, t)$  if and only if  $x_1 = t$ .

- (35) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $R$  be an order sorted congruence of  $\text{ParsedTermsOSAX}$ , and  $t_2, t_3$  be elements of  $\text{TS}(\text{DTConOSAX})$ . Then  $t_3 \in \text{OSClass}(R, t_2)$  if and only if  $\text{OSClass}(R, t_2) = \text{OSClass}(R, t_3)$ .

- (36) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $R_1, R_2$  be order sorted congruences of  $\text{ParsedTermsOSAX}$ , and  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ . If  $R_1 \subseteq R_2$ , then  $\text{OSClass}(R_1, t) \subseteq \text{OSClass}(R_2, t)$ .

- (37) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $s$  be an element of  $S$ ,  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ , and  $x, x_1$  be sets. Suppose  $x \in X(s)$  and  $t$  = the root tree of  $\langle x, s \rangle$ . Then  $x_1 \in \text{OSClass}(\text{LCongruenceX}, t)$  if and only if  $x_1 = t$ .



Let  $S$  be a locally directed order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , let  $A$  be a non-empty many sorted set indexed by the carrier of  $S$ , let  $F$  be a many sorted function from  $\text{PTVars}X$  into  $A$ , and let  $t$  be a symbol of  $\text{DTConOSAX}$ . Let us assume that  $t \in$  the terminals of  $\text{DTConOSAX}$ . The functor  $\pi(F, A, t)$  yielding an element of  $\bigcup A$  is defined as follows:

(Def. 29) For every function  $f$  such that  $f = F(t_2)$  holds  $\pi(F, A, t) = f(\text{the root tree of } t)$ .

One can prove the following proposition

(38) Let  $S$  be a locally directed order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $U_1$  be a monotone non-empty order sorted algebra of  $S$ , and  $f$  be a many sorted function from  $\text{PTVars}X$  into the sorts of  $U_1$ . Then there exists a many sorted function  $h$  from  $\text{ParsedTermsOSAX}$  into  $U_1$  such that  $h$  is a homomorphism of  $\text{ParsedTermsOSAX}$  into  $U_1$  and order-sorted and  $h \upharpoonright \text{PTVars}X = f$ .

Let  $S$  be a locally directed order sorted signature, let  $X$  be a non-empty many sorted set indexed by  $S$ , and let  $s$  be an element of  $S$ . The functor  $\text{NHReverse}(s, X)$  yields a function from  $\text{OSFreeGen}(s, X)$  into  $\text{PTVars}(s, X)$  and is defined by the condition (Def. 30).

(Def. 30) Let  $t$  be a symbol of  $\text{DTConOSAX}$ . Suppose  $(\text{OSNatHom}(\text{ParsedTermsOSAX}, \text{LCongruence}X))(s)(\text{the root tree of } t) \in \text{OSFreeGen}(s, X)$ . Then  $(\text{NHReverse}(s, X))((\text{OSNatHom}(\text{ParsedTermsOSAX}, \text{LCongruence}X))(s)(\text{the root tree of } t)) = \text{the root tree of } t$ .

Let  $S$  be a locally directed order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $\text{NHReverse}X$  yields a many sorted function from  $\text{OSFreeGen}X$  into  $\text{PTVars}X$  and is defined by:

(Def. 31) For every element  $s$  of  $S$  holds  $(\text{NHReverse}X)(s) = \text{NHReverse}(s, X)$ .

We now state two propositions:

(39) Let  $S$  be a locally directed order sorted signature and  $X$  be a non-empty many sorted set indexed by  $S$ . Then  $\text{OSFreeGen}X$  is *osfree*.

(40) Let  $S$  be a locally directed order sorted signature and  $X$  be a non-empty many sorted set indexed by  $S$ . Then  $\text{FreeOSAX}$  is *osfree*.

Let  $S$  be a locally directed order sorted signature. Observe that there exists a non-empty monotone order sorted algebra of  $S$  which is *osfree* and *strict*.

### 3. MINIMAL TERMS

Let  $S$  be a locally directed regular monotone order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $\text{PTMin}X$  yields a function from  $\text{TS}(\text{DTConOSAX})$  into  $\text{TS}(\text{DTConOSAX})$  and is defined by the conditions (Def. 32).

(Def. 32)(i) For every symbol  $t$  of  $\text{DTConOSAX}$  such that  $t \in$  the terminals of  $\text{DTConOSAX}$  holds  $(\text{PTMin}X)(\text{the root tree of } t) = \prod t$ , and

(ii) for every symbol  $n_1$  of  $\text{DTConOSAX}$  and for every finite sequence  $t_1$  of elements of  $\text{TS}(\text{DTConOSAX})$  such that  $n_1 \Rightarrow$  the roots of  $t_1$  holds  $(\text{PTMin}X)(n_1\text{-tree}(t_1)) = \pi_{\text{PTMin}X \cdot t_1}(@ (X, n_1))$ .

The following propositions are true:

(41) Let  $S$  be a locally directed regular monotone order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $t$  be an element of  $\text{TS}(\text{DTConOSAX})$ . Then

(i)  $(\text{PTMin}X)(t) \in \text{OSClass}(\text{PTCongruence}X, t)$ ,

(ii)  $\text{LeastSort}(\text{PTMin}X)(t) \leq \text{LeastSort}t$ ,

- (iii) for every element  $s$  of  $S$  and for every set  $x$  such that  $x \in X(s)$  and  $t =$  the root tree of  $\langle x, s \rangle$  holds  $(PTMinX)(t) = t$ , and
- (iv) for every operation symbol  $o$  of  $S$  and for every finite sequence  $t_1$  of elements of  $TS(DTConOSAX)$  such that  $OSSym(o, X) \Rightarrow$  the roots of  $t_1$  and  $t = OSSym(o, X)$ -tree( $t_1$ ) holds  $LeastSortsPTMinX \cdot t_1 \leq Arity(o)$  and  $OSSym(o, X) \Rightarrow$  the roots of  $PTMinX \cdot t_1$  and  $OSSym(LBound(o, LeastSortsPTMinX \cdot t_1), X) \Rightarrow$  the roots of  $PTMinX \cdot t_1$  and  $(PTMinX)(t) = OSSym(LBound(o, LeastSortsPTMinX \cdot t_1), X)$ -tree( $PTMinX \cdot t_1$ ).
- (42) Let  $S$  be a locally directed regular monotone order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $t, t_2$  be elements of  $TS(DTConOSAX)$ . If  $t_2 \in OSClass(PTCongruenceX, t)$ , then  $(PTMinX)(t_2) = (PTMinX)(t)$ .
- (43) Let  $S$  be a locally directed regular monotone order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $t_2, t_3$  be elements of  $TS(DTConOSAX)$ . Then  $t_3 \in OSClass(PTCongruenceX, t_2)$  if and only if  $(PTMinX)(t_3) = (PTMinX)(t_2)$ .
- (44) Let  $S$  be a locally directed regular monotone order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $t_2$  be an element of  $TS(DTConOSAX)$ . Then  $(PTMinX)((PTMinX)(t_2)) = (PTMinX)(t_2)$ .
- (45) Let  $S$  be a locally directed regular monotone order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ ,  $R$  be a monotone equivalence order sorted relation of  $ParsedTermsOSAX$ , and  $t$  be an element of  $TS(DTConOSAX)$ . Then  $\langle t, (PTMinX)(t) \rangle \in R(LeastSortt)$ .
- (46) Let  $S$  be a locally directed regular monotone order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $R$  be a monotone equivalence order sorted relation of  $ParsedTermsOSAX$ . Then  $PTCongruenceX \subseteq R$ .
- (47) Let  $S$  be a locally directed regular monotone order sorted signature and  $X$  be a non-empty many sorted set indexed by  $S$ . Then  $LCongruenceX = PCongruenceX$ .

Let  $S$  be a locally directed regular monotone order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . An element of  $TS(DTConOSAX)$  is called a minimal term of  $S, X$  if:

(Def. 33)  $(PTMinX)(it) = it$ .

Let  $S$  be a locally directed regular monotone order sorted signature and let  $X$  be a non-empty many sorted set indexed by  $S$ . The functor  $MinTermsX$  yielding a subset of  $TS(DTConOSAX)$  is defined by:

(Def. 34)  $MinTermsX = rng PTMinX$ .

The following proposition is true

- (48) Let  $S$  be a locally directed regular monotone order sorted signature,  $X$  be a non-empty many sorted set indexed by  $S$ , and  $x$  be a set. Then  $x$  is a minimal term of  $S, X$  if and only if  $x \in MinTermsX$ .

#### ACKNOWLEDGMENTS

Thanks to Joseph Goguen, for providing me with his articles on osas, and Andrzej Trybulec, for suggesting and funding this work in Bialystok.

## REFERENCES

- [1] Grzegorz Bancerek. Curried and uncurried functions. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol12/funct\\_5.html](http://mizar.org/JFM/Vol12/funct_5.html).
- [2] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol12/card\\_3.html](http://mizar.org/JFM/Vol12/card_3.html).
- [3] Grzegorz Bancerek. Cartesian product of functions. *Journal of Formalized Mathematics*, 3, 1991. [http://mizar.org/JFM/Vol13/funct\\_6.html](http://mizar.org/JFM/Vol13/funct_6.html).
- [4] Grzegorz Bancerek. König's Lemma. *Journal of Formalized Mathematics*, 3, 1991. [http://mizar.org/JFM/Vol13/trees\\_2.html](http://mizar.org/JFM/Vol13/trees_2.html).
- [5] Grzegorz Bancerek. Sets and functions of trees and joining operations of trees. *Journal of Formalized Mathematics*, 4, 1992. [http://mizar.org/JFM/Vol14/trees\\_3.html](http://mizar.org/JFM/Vol14/trees_3.html).
- [6] Grzegorz Bancerek. Joining of decorated trees. *Journal of Formalized Mathematics*, 5, 1993. [http://mizar.org/JFM/Vol15/trees\\_4.html](http://mizar.org/JFM/Vol15/trees_4.html).
- [7] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol11/finseq\\_1.html](http://mizar.org/JFM/Vol11/finseq_1.html).
- [8] Grzegorz Bancerek and Piotr Rudnicki. On defining functions on trees. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol15/dtconstr.html>.
- [9] Ewa Burakowska. Subalgebras of many sorted algebra. Lattice of subalgebras. *Journal of Formalized Mathematics*, 6, 1994. [http://mizar.org/JFM/Vol16/msualg\\_2.html](http://mizar.org/JFM/Vol16/msualg_2.html).
- [10] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol11/funct\\_1.html](http://mizar.org/JFM/Vol11/funct_1.html).
- [11] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol11/funct\\_2.html](http://mizar.org/JFM/Vol11/funct_2.html).
- [12] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol11/partfun1.html>.
- [13] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol11/zfmisc\\_1.html](http://mizar.org/JFM/Vol11/zfmisc_1.html).
- [14] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol12/finseq\\_2.html](http://mizar.org/JFM/Vol12/finseq_2.html).
- [15] Patricia L. Carlson and Grzegorz Bancerek. Context-free grammar — part I. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol14/lang1.html>.
- [16] Małgorzata Korolkiewicz. Homomorphisms of many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. [http://mizar.org/JFM/Vol16/msualg\\_3.html](http://mizar.org/JFM/Vol16/msualg_3.html).
- [17] Małgorzata Korolkiewicz. Many sorted quotient algebra. *Journal of Formalized Mathematics*, 6, 1994. [http://mizar.org/JFM/Vol16/msualg\\_4.html](http://mizar.org/JFM/Vol16/msualg_4.html).
- [18] Beata Madras. Product of family of universal algebras. *Journal of Formalized Mathematics*, 5, 1993. [http://mizar.org/JFM/Vol15/pralg\\_1.html](http://mizar.org/JFM/Vol15/pralg_1.html).
- [19] Andrzej Nędzusiak.  $\sigma$ -fields and probability. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol11/prob\\_1.html](http://mizar.org/JFM/Vol11/prob_1.html).
- [20] Beata Perkowska. Free many sorted universal algebra. *Journal of Formalized Mathematics*, 6, 1994. <http://mizar.org/JFM/Vol16/msafree.html>.
- [21] Konrad Raczkowski and Paweł Sadowski. Equivalence relations and classes of abstraction. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol11/eqrel\\_1.html](http://mizar.org/JFM/Vol11/eqrel_1.html).
- [22] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [23] Andrzej Trybulec. Tuples, projections and Cartesian products. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol11/mcart\\_1.html](http://mizar.org/JFM/Vol11/mcart_1.html).
- [24] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol15/pboole.html>.
- [25] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. [http://mizar.org/JFM/Vol16/msualg\\_1.html](http://mizar.org/JFM/Vol16/msualg_1.html).
- [26] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol11/orders\\_1.html](http://mizar.org/JFM/Vol11/orders_1.html).
- [27] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol12/finseq\\_4.html](http://mizar.org/JFM/Vol12/finseq_4.html).
- [28] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol11/subset\\_1.html](http://mizar.org/JFM/Vol11/subset_1.html).
- [29] Josef Urban. Homomorphisms of order sorted algebras. *Journal of Formalized Mathematics*, 14, 2002. [http://mizar.org/JFM/Vol14/osalg\\_3.html](http://mizar.org/JFM/Vol14/osalg_3.html).

- [30] Josef Urban. Order sorted algebras. *Journal of Formalized Mathematics*, 14, 2002. [http://mizar.org/JFM/Voll4/osalg\\_1.html](http://mizar.org/JFM/Voll4/osalg_1.html).
- [31] Josef Urban. Order sorted quotient algebra. *Journal of Formalized Mathematics*, 14, 2002. [http://mizar.org/JFM/Voll4/osalg\\_4.html](http://mizar.org/JFM/Voll4/osalg_4.html).
- [32] Josef Urban. Subalgebras of a order sorted algebra. Lattice of subalgebras. *Journal of Formalized Mathematics*, 14, 2002. [http://mizar.org/JFM/Voll4/osalg\\_2.html](http://mizar.org/JFM/Voll4/osalg_2.html).
- [33] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/relat\\_1.html](http://mizar.org/JFM/Voll/relat_1.html).
- [34] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/relset\\_1.html](http://mizar.org/JFM/Voll/relset_1.html).
- [35] Edmund Woronowicz and Anna Zalewska. Properties of binary relations. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/relat\\_2.html](http://mizar.org/JFM/Voll/relat_2.html).

*Received September 19, 2002*

*Published January 2, 2004*

---