

Construction of a bilinear antisymmetric form in symplectic vector space¹

Eugeniusz Kusak
Warsaw University
Białystok

Wojciech Leończuk
Warsaw University
Białystok

Michał Muzalewski
Warsaw University
Białystok

Summary. In this text we will present unpublished results by Eugeniusz Kusak. It contains an axiomatic description of the class of all spaces $\langle V; \perp_\xi \rangle$, where V is a vector space over a field F , $\xi : V \times V \rightarrow F$ is a bilinear antisymmetric form i.e. $\xi(x,y) = -\xi(y,x)$ and $x \perp_\xi y$ iff $\xi(x,y) = 0$ for $x, y \in V$. It also contains an effective construction of bilinear antisymmetric form ξ for given symplectic space $\langle V; \perp \rangle$ such that $\perp = \perp_\xi$. The basic tool used in this method is the notion of orthogonal projection $J(a,b,x)$ for $a, b, x \in V$. We should stress the fact that axioms of orthogonal and symplectic spaces differ only by one axiom, namely: $x \perp y + \varepsilon z \& y \perp z + \varepsilon x \Rightarrow z \perp x + \varepsilon y$. For $\varepsilon = +1$ we get the axiom characterizing symplectic geometry. For $\varepsilon = -1$ we get the axiom on three perpendiculars characterizing orthogonal geometry - see [5].

MML Identifier: SYMSP_1.

WWW: http://mizar.org/JFM/Vol1/symsp_1.html

The articles [6], [3], [8], [1], [2], [7], [4], and [9] provide the notation and terminology for this paper.

In this paper F denotes a field.

Let us consider F . We consider symplectic structures over F as extensions of vector space structure over F as systems

\langle a carrier, an addition, a zero, a left multiplication, an orthogonality \rangle ,
where the carrier is a set, the addition is a binary operation on the carrier, the zero is an element of the carrier, the left multiplication is a function from [the carrier of F , the carrier] into the carrier, and the orthogonality is a binary relation on the carrier.

Let us consider F . Note that there exists a symplectic structure over F which is non empty.

Let us consider F , let S be a symplectic structure over F , and let a, b be elements of S . The predicate $a \perp b$ is defined as follows:

(Def. 1) $\langle a, b \rangle \in$ the orthogonality of S .

Let us consider F , let X be a non empty set, let m_1 be a binary operation on X , let o be an element of X , let m_2 be a function from [the carrier of F , X] into X , and let m_3 be a binary relation on X . Observe that $\langle X, m_1, o, m_2, m_3 \rangle$ is non empty.

Let us consider F . One can verify that there exists a non empty symplectic structure over F which is Abelian, add-associative, right zeroed, and right complementable.

Let us consider F and let I_1 be an Abelian add-associative right zeroed right complementable non empty symplectic structure over F . We say that I_1 is symplectic space-like if and only if the condition (Def. 2) is satisfied.

¹Supported by RPBP.III-24.C6.

(Def. 2) Let a, b, c, x be elements of I_1 and l be an element of F . Then

- (i) if $a \neq 0_{(I_1)}$, then there exists an element y of the carrier of I_1 such that $y \not\perp a$,
- (ii) if $a \perp b$, then $l \cdot a \perp b$,
- (iii) if $b \perp a$ and $c \perp a$, then $b + c \perp a$,
- (iv) if $b \not\perp a$, then there exists an element k of F such that $x - k \cdot b \perp a$, and
- (v) if $a \perp b + c$ and $b \perp c + a$, then $c \perp a + b$.

Let us consider F . Note that there exists an Abelian add-associative right zeroed right complementable non empty simplectic structure over F which is simplectic space-like, vector space-like, and strict.

Let us consider F . A simplectic space over F is a simplectic space-like vector space-like Abelian add-associative right zeroed right complementable non empty simplectic structure over F .

We adopt the following rules: S denotes a simplectic space over F , $a, b, c, d, a', b', p, q, x, y, z$ denote elements of S , and k, l denote elements of F .

One can prove the following propositions:

- (11)¹ $0_S \perp a$.
- (12) If $a \perp b$, then $b \perp a$.
- (13) If $a \not\perp b$ and $c + a \perp b$, then $c \not\perp b$.
- (14) If $b \not\perp a$ and $c \perp a$, then $b + c \not\perp a$.
- (15) If $b \not\perp a$ and $l \neq 0_F$, then $l \cdot b \not\perp a$ and $b \not\perp l \cdot a$.
- (16) If $a \perp b$, then $-a \perp b$.
- (19)² If $a \not\perp c$, then $a + b \not\perp c$ or $(\mathbf{1}_F + \mathbf{1}_F) \cdot a + b \not\perp c$.
- (20) If $a' \not\perp a$ and $a' \perp b$ and $b' \not\perp b$ and $b' \perp a$, then $a' + b' \not\perp a$ and $a' + b' \not\perp b$.
- (21) If $a \neq 0_S$ and $b \neq 0_S$, then there exists p such that $p \not\perp a$ and $p \not\perp b$.
- (22) If $\mathbf{1}_F + \mathbf{1}_F \neq 0_F$ and $a \neq 0_S$ and $b \neq 0_S$ and $c \neq 0_S$, then there exists p such that $p \not\perp a$ and $p \not\perp b$ and $p \not\perp c$.
- (23) If $a - b \perp d$ and $a - c \perp d$, then $b - c \perp d$.
- (24) If $b \not\perp a$ and $x - k \cdot b \perp a$ and $x - l \cdot b \perp a$, then $k = l$.
- (25) If $\mathbf{1}_F + \mathbf{1}_F \neq 0_F$, then $a \perp a$.

Let us consider F and let us consider S, a, b, x . Let us assume that $b \not\perp a$. The functor $J(a, b, x)$ yielding an element of F is defined by:

(Def. 6)³ For every element l of F such that $x - l \cdot b \perp a$ holds $J(a, b, x) = l$.

The following propositions are true:

- (27)⁴ If $b \not\perp a$, then $x - J(a, b, x) \cdot b \perp a$.
- (28) If $b \not\perp a$, then $J(a, b, l \cdot x) = l \cdot J(a, b, x)$.
- (29) If $b \not\perp a$, then $J(a, b, x + y) = J(a, b, x) + J(a, b, y)$.
- (30) If $b \not\perp a$ and $l \neq 0_F$, then $J(a, l \cdot b, x) = l^{-1} \cdot J(a, b, x)$.

¹ The propositions (1)–(10) have been removed.

² The propositions (17) and (18) have been removed.

³ The definitions (Def. 3)–(Def. 5) have been removed.

⁴ The proposition (26) has been removed.

- (31) If $b \not\perp a$ and $l \neq 0_F$, then $J(l \cdot a, b, x) = J(a, b, x)$.
- (32) If $b \not\perp a$ and $p \perp a$, then $J(a, b + p, c) = J(a, b, c)$ and $J(a, b, c + p) = J(a, b, c)$.
- (33) If $b \not\perp a$ and $p \perp b$ and $p \perp c$, then $J(a + p, b, c) = J(a, b, c)$.
- (34) If $b \not\perp a$ and $c - b \perp a$, then $J(a, b, c) = \mathbf{1}_F$.
- (35) If $b \not\perp a$, then $J(a, b, b) = \mathbf{1}_F$.
- (36) If $b \not\perp a$, then $x \perp a$ iff $J(a, b, x) = 0_F$.
- (37) If $b \not\perp a$ and $q \not\perp a$, then $J(a, b, p) \cdot J(a, b, q)^{-1} = J(a, q, p)$.
- (38) If $b \not\perp a$ and $c \not\perp a$, then $J(a, b, c) = J(a, c, b)^{-1}$.
- (39) If $b \not\perp a$ and $b \perp c + a$, then $J(a, b, c) = J(c, b, a)$.
- (40) If $a \not\perp b$ and $c \not\perp b$, then $J(c, b, a) = (-J(b, a, c)^{-1}) \cdot J(a, b, c)$.
- (41) If $\mathbf{1}_F + \mathbf{1}_F \neq 0_F$ and $a \not\perp p$ and $a \not\perp q$ and $b \not\perp p$ and $b \not\perp q$, then $J(a, p, q) \cdot J(b, q, p) = J(p, a, b) \cdot J(q, b, a)$.
- (42) If $\mathbf{1}_F + \mathbf{1}_F \neq 0_F$ and $p \not\perp a$ and $p \not\perp x$ and $q \not\perp a$ and $q \not\perp x$, then $J(a, q, p) \cdot J(p, a, x) = J(x, q, p) \cdot J(q, a, x)$.
- (43) If $\mathbf{1}_F + \mathbf{1}_F \neq 0_F$ and $p \not\perp a$ and $p \not\perp x$ and $q \not\perp a$ and $q \not\perp x$ and $b \not\perp a$, then $J(a, b, p) \cdot J(p, a, x) \cdot J(x, p, y) = J(a, b, q) \cdot J(q, a, x) \cdot J(x, q, y)$.
- (44) If $a \not\perp p$ and $x \not\perp p$ and $y \not\perp p$, then $J(p, a, x) \cdot J(x, p, y) = (-J(p, a, y)) \cdot J(y, p, x)$.

Let us consider F, S, x, y, a, b . Let us assume that $b \not\perp a$ and $\mathbf{1}_F + \mathbf{1}_F \neq 0_F$. The functor $x \cdot_{a,b} y$ yielding an element of F is defined by:

- (Def. 7)(i) For every q such that $q \not\perp a$ and $q \not\perp x$ holds $x \cdot_{a,b} y = J(a, b, q) \cdot J(q, a, x) \cdot J(x, q, y)$ if there exists p such that $p \not\perp a$ and $p \not\perp x$,
- (ii) $x \cdot_{a,b} y = 0_F$ if for every p holds $p \perp a$ or $p \perp x$.

We now state several propositions:

- (47)⁵ If $\mathbf{1}_F + \mathbf{1}_F \neq 0_F$ and $b \not\perp a$ and $x = 0_S$, then $x \cdot_{a,b} y = 0_F$.
- (48) If $\mathbf{1}_F + \mathbf{1}_F \neq 0_F$ and $b \not\perp a$, then $x \cdot_{a,b} y = 0_F$ iff $y \perp x$.
- (49) If $\mathbf{1}_F + \mathbf{1}_F \neq 0_F$ and $b \not\perp a$, then $x \cdot_{a,b} y = -y \cdot_{a,b} x$.
- (50) If $\mathbf{1}_F + \mathbf{1}_F \neq 0_F$ and $b \not\perp a$, then $x \cdot_{a,b} (l \cdot y) = l \cdot x \cdot_{a,b} y$.
- (51) If $\mathbf{1}_F + \mathbf{1}_F \neq 0_F$ and $b \not\perp a$, then $x \cdot_{a,b} (y + z) = x \cdot_{a,b} y + x \cdot_{a,b} z$.

REFERENCES

- [1] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [2] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [3] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [4] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/vectsp_1.html.
- [5] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Construction of a bilinear symmetric form in orthogonal vector space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ortsp_1.html.

⁵ The propositions (45) and (46) have been removed.

- [6] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [7] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/rvect_1.html.
- [8] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [9] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relset_1.html.

Received November 23, 1989

Published January 2, 2004
