

Injective Spaces¹

Jarosław Gryko
University of Białystok

MML Identifier: WAYBEL18.

WWW: <http://mizar.org/JFM/Vol10/waybel18.html>

The articles [23], [22], [11], [28], [19], [29], [31], [9], [10], [20], [8], [2], [7], [25], [1], [17], [27], [15], [18], [21], [13], [14], [12], [24], [3], [30], [6], [4], [5], [16], and [26] provide the notation and terminology for this paper.

1. PRODUCT TOPOLOGIES

One can prove the following propositions:

- (1) Let x, y, z, Z be sets. Then $Z \subseteq \{x, y, z\}$ if and only if one of the following conditions is satisfied:
 $Z = \emptyset$ or $Z = \{x\}$ or $Z = \{y\}$ or $Z = \{z\}$ or $Z = \{x, y\}$ or $Z = \{y, z\}$ or $Z = \{x, z\}$ or $Z = \{x, y, z\}$.
- (2) For every set X and for all families A, B of subsets of X such that $B = A \setminus \{\emptyset\}$ or $A = B \cup \{\emptyset\}$ holds $\text{UniCl}(A) = \text{UniCl}(B)$.
- (3) Let T be a topological space and K be a family of subsets of T . Then K is a basis of T if and only if $K \setminus \{\emptyset\}$ is a basis of T .

Let F be a binary relation. We say that F is topological space yielding if and only if:

(Def. 1) For every set x such that $x \in \text{rng } F$ holds x is a topological structure.

Let us note that every function which is topological space yielding is also 1-sorted yielding.

Let I be a set. One can check that there exists a many sorted set indexed by I which is topological space yielding.

Let I be a set. Observe that there exists a many sorted set indexed by I which is topological space yielding and nonempty.

Let J be a non empty set, let A be a topological space yielding many sorted set indexed by J , and let j be an element of J . Then $A(j)$ is a topological structure.

Let I be a set and let J be a topological space yielding many sorted set indexed by I . The product prebasis for J is a family of subsets of \prod (the support of J) and is defined by the condition (Def. 2).

(Def. 2) Let x be a subset of \prod (the support of J). Then $x \in$ the product prebasis for J if and only if there exists a set i and there exists a topological structure T and there exists a subset V of T such that $i \in I$ and V is open and $T = J(i)$ and $x = \prod$ ((the support of J) $\dot{+}$ (i, V)).

Next we state the proposition

¹This work has been supported by KBN Grant 8 T11C 018 12.

- (4) For every set X and for every family A of subsets of X holds $\langle X, \text{UniCl}(\text{FinMeetCl}(A)) \rangle$ is topological space-like.

Let I be a set and let J be a topological space yielding nonempty many sorted set indexed by I . The functor $\prod J$ yields a strict topological space and is defined by:

- (Def. 3) The carrier of $\prod J = \prod(\text{the support of } J)$ and the product prebasis for J is a prebasis of $\prod J$.

Let I be a set and let J be a topological space yielding nonempty many sorted set indexed by I . One can check that $\prod J$ is non empty.

Let I be a non empty set, let J be a topological space yielding nonempty many sorted set indexed by I , and let i be an element of I . Then $J(i)$ is a non empty topological structure.

Let I be a set and let J be a topological space yielding nonempty many sorted set indexed by I . One can check that every element of $\prod J$ is function-like and relation-like.

Let I be a non empty set, let J be a topological space yielding nonempty many sorted set indexed by I , let x be an element of $\prod J$, and let i be an element of I . Then $x(i)$ is an element of $J(i)$.

Let I be a non empty set, let J be a topological space yielding nonempty many sorted set indexed by I , and let i be an element of I . The functor $\text{proj}(J, i)$ yields a map from $\prod J$ into $J(i)$ and is defined by:

- (Def. 4) $\text{proj}(J, i) = \text{proj}(\text{the support of } J, i)$.

Next we state three propositions:

- (5) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I , i be an element of I , and P be a subset of $J(i)$. Then $(\text{proj}(J, i))^{-1}(P) = \prod(\text{the support of } J) + \cdot (i, P)$.
- (6) Let I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I , and i be an element of I . Then $\text{proj}(J, i)$ is continuous.
- (7) Let X be a non empty topological space, I be a non empty set, J be a topological space yielding nonempty many sorted set indexed by I , and f be a map from X into $\prod J$. Then f is continuous if and only if for every element i of I holds $\text{proj}(J, i) \cdot f$ is continuous.

2. INJECTIVE SPACES

Let Z be a topological structure. We say that Z is injective if and only if the condition (Def. 5) is satisfied.

- (Def. 5) Let X be a non empty topological space and f be a map from X into Z . Suppose f is continuous. Let Y be a non empty topological space. Suppose X is a subspace of Y . Then there exists a map g from Y into Z such that g is continuous and $g|_{\text{the carrier of } X} = f$.

The following two propositions are true:

- (8) Let I be a non empty set and J be a topological space yielding nonempty many sorted set indexed by I . If for every element i of I holds $J(i)$ is injective, then $\prod J$ is injective.
- (9) Let T be a non empty topological space. Suppose T is injective. Let S be a non empty subspace of T . If S is a retract of T , then S is injective.

Let X be a 1-sorted structure, let Y be a topological structure, and let f be a map from X into Y . The functor $\text{Im } f$ yielding a subspace of Y is defined by:

- (Def. 6) $\text{Im } f = Y \upharpoonright \text{rng } f$.

Let X be a non empty 1-sorted structure, let Y be a non empty topological structure, and let f be a map from X into Y . One can verify that $\text{Im } f$ is non empty.

Next we state the proposition

- (10) Let X be a 1-sorted structure, Y be a topological structure, and f be a map from X into Y . Then the carrier of $\text{Im } f = \text{rng } f$.

Let X be a 1-sorted structure, let Y be a non empty topological structure, and let f be a map from X into Y . The functor f° yielding a map from X into $\text{Im } f$ is defined by:

(Def. 7) $f^\circ = f$.

One can prove the following proposition

- (11) Let X, Y be non empty topological spaces and f be a map from X into Y . If f is continuous, then f° is continuous.

Let X be a 1-sorted structure, let Y be a non empty topological structure, and let f be a map from X into Y . Observe that f° is onto.

Let X, Y be topological structures. We say that X is a topological retract of Y if and only if:

(Def. 8) There exists a map f from Y into Y such that f is continuous and $f \cdot f = f$ and $\text{Im } f$ and X are homeomorphic.

Next we state the proposition

- (12) Let T, S be non empty topological spaces. Suppose T is injective. Let f be a map from T into S . If f° is a homeomorphism, then T is a topological retract of S .

The Sierpiński space is a strict topological structure and is defined by the conditions (Def. 9).

(Def. 9)(i) The carrier of the Sierpiński space = $\{0, 1\}$, and

(ii) the topology of the Sierpiński space = $\{\emptyset, \{1\}, \{0, 1\}\}$.

One can check that the Sierpiński space is non empty and topological space-like.

Let us mention that the Sierpiński space is discernible.

Let us note that the Sierpiński space is injective.

Let I be a set and let S be a non empty 1-sorted structure. One can verify that $I \mapsto S$ is nonempty.

Let I be a set and let T be a topological structure. Observe that $I \mapsto T$ is topological space yielding.

Let I be a set and let L be a reflexive relational structure. One can verify that $I \mapsto L$ is reflexive-yielding.

Let I be a non empty set and let L be a non empty antisymmetric relational structure. Note that $\prod(I \mapsto L)$ is antisymmetric.

Let I be a non empty set and let L be a non empty transitive relational structure. Note that $\prod(I \mapsto L)$ is transitive.

We now state two propositions:

- (13) Let T be a Scott topological augmentation of $2_{\underline{\mathbb{C}}}^1$. Then the topology of T = the topology of the Sierpiński space.

- (14) Let I be a non empty set. Then $\{\prod((\text{the support of } I \mapsto \text{the Sierpiński space}) + \cdot (i, \{1\})) : i \text{ ranges over elements of } I\}$ is a prebasis of $\prod(I \mapsto \text{the Sierpiński space})$.

Let I be a non empty set and let L be a complete lattice. Note that $\prod(I \mapsto L)$ is complete and has l.u.b.'s.

Let I be a non empty set and let X be an algebraic lower-bounded lattice. Note that $\prod(I \mapsto X)$ is algebraic.

We now state several propositions:

- (15) Let X be a non empty set. Then there exists a map f from $2_{\underline{\mathbb{C}}}^X$ into $\prod(X \mapsto 2_{\underline{\mathbb{C}}}^1)$ such that f is isomorphic and for every subset Y of X holds $f(Y) = \chi_{Y,X}$.

- (16) Let I be a non empty set and T be a Scott topological augmentation of $\prod(I \mapsto 2_{\subseteq}^1)$. Then the topology of $T =$ the topology of $\prod(I \mapsto$ the Sierpiński space).
- (17) Let T, S be non empty topological spaces. Suppose the carrier of $T =$ the carrier of S and the topology of $T =$ the topology of S and T is injective. Then S is injective.
- (18) For every non empty set I holds every Scott topological augmentation of $\prod I \mapsto 2_{\subseteq}^1$ is injective.
- (19) Let T be a T_0 -space. Then there exists a non empty set M and there exists a map f from T into $\prod(M \mapsto$ the Sierpiński space) such that f° is a homeomorphism.
- (20) Let T be a T_0 -space. Suppose T is injective. Then there exists a non empty set M such that T is a topological retract of $\prod(M \mapsto$ the Sierpiński space).

REFERENCES

- [1] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.
- [2] Grzegorz Bancerek. Cartesian product of functions. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol13/funct_6.html.
- [3] Grzegorz Bancerek. Bounds in posets and relational substructures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_0.html.
- [4] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_0.html.
- [5] Grzegorz Bancerek. The "way-below" relation. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_3.html.
- [6] Grzegorz Bancerek. Bases and refinements of topologies. *Journal of Formalized Mathematics*, 10, 1998. http://mizar.org/JFM/Vol10/yellow_9.html.
- [7] Grzegorz Bancerek and Andrzej Trybulec. Miscellaneous facts about functions. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/funct_7.html.
- [8] Czesław Byliński. Basic functions and operations on functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_3.html.
- [9] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [10] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [11] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [12] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/tops_2.html.
- [13] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.
- [14] Mariusz Giero. More on products of many sorted algebras. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/pralg_3.html.
- [15] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_1.html.
- [16] Artur Kornilowicz. On the topological properties of meet-continuous lattices. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_9.html.
- [17] Beata Madras. Product of family of universal algebras. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/pralg_1.html.
- [18] Yatsuka Nakamura, Piotr Rudnicki, Andrzej Trybulec, and Pauline N. Kawamoto. Preliminaries to circuits, I. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/pre_circ.html.
- [19] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/setfam_1.html.
- [20] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.
- [21] Alexander Yu. Shibakov and Andrzej Trybulec. The Cantor set. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/cantor_1.html.
- [22] Andrzej Trybulec. Enumerated sets. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/enumset1.html>.

- [23] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [24] Andrzej Trybulec. A Borsuk theorem on homotopy types. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/borsuk_1.html.
- [25] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/pboole.html>.
- [26] Andrzej Trybulec. Scott topology. *Journal of Formalized Mathematics*, 9, 1997. <http://mizar.org/JFM/Vol9/waybell1.html>.
- [27] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/orders_1.html.
- [28] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [29] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.
- [30] Mariusz Żynel and Czesław Byliński. Properties of relational structures, posets, lattices and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_2.html.
- [31] Mariusz Żynel and Adam Guzowski. T_0 topological spaces. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/t_toposp.html.

Received April 17, 1998

Published January 2, 2004
