

On the Characterization of Modular and Distributive Lattices¹

Adam Naumowicz
University of Białystok

Summary. This article contains definitions of the “pentagon” lattice N_5 and the “diamond” lattice M_3 . It is followed by the characterization of modular and distributive lattices depending on the possible shape of substructures. The last part treats of interval-like sublattices of any lattice.

MML Identifier: YELLOW11.

WWW: <http://mizar.org/JFM/Vol10/yellow11.html>

The articles [11], [10], [13], [14], [5], [6], [12], [2], [3], [9], [15], [7], [8], [4], and [1] provide the notation and terminology for this paper.

1. PRELIMINARIES

The following propositions are true:

- (1) $3 = \{0, 1, 2\}$.
- (2) $2 \setminus 1 = \{1\}$.
- (3) $3 \setminus 1 = \{1, 2\}$.
- (4) $3 \setminus 2 = \{2\}$.

2. MAIN PART

One can prove the following propositions:

- (5) Let L be an antisymmetric reflexive relational structure with g.l.b.'s and l.u.b.'s and a, b be elements of L . Then $a \sqcap b = b$ if and only if $a \sqcup b = a$.
- (6) For every lattice L and for all elements a, b, c of L holds $(a \sqcap b) \sqcup (a \sqcap c) \leq a \sqcap (b \sqcup c)$.
- (7) For every lattice L and for all elements a, b, c of L holds $a \sqcup (b \sqcap c) \leq (a \sqcup b) \sqcap (a \sqcup c)$.
- (8) For every lattice L and for all elements a, b, c of L such that $a \leq c$ holds $a \sqcup (b \sqcap c) \leq (a \sqcup b) \sqcap c$.

The relational structure N_5 is defined as follows:

(Def. 1) $N_5 = \langle \{0, 3 \setminus 1, 2, 3 \setminus 2, 3\}, \subseteq \rangle$.

¹This work has been supported by KBN Grant 8 T11C 018 12.

Let us note that N_5 is strict, reflexive, transitive, and antisymmetric and N_5 has g.l.b.'s and l.u.b.'s. The relational structure M_3 is defined by:

(Def. 2) $M_3 = \langle \{0, 1, 2 \setminus 1, 3 \setminus 2, 3\}, \subseteq \rangle$.

Let us note that M_3 is strict, reflexive, transitive, and antisymmetric and M_3 has g.l.b.'s and l.u.b.'s.

One can prove the following propositions:

- (9) Let L be a lattice. Then the following statements are equivalent
- (i) there exists a full sublattice K of L such that N_5 and K are isomorphic,
 - (ii) there exist elements a, b, c, d, e of L such that $a \neq b$ and $a \neq c$ and $a \neq d$ and $a \neq e$ and $b \neq c$ and $b \neq d$ and $b \neq e$ and $c \neq d$ and $c \neq e$ and $d \neq e$ and $a \sqcap b = a$ and $a \sqcap c = a$ and $c \sqcap e = c$ and $d \sqcap e = d$ and $b \sqcap c = a$ and $b \sqcap d = b$ and $c \sqcap d = a$ and $b \sqcup c = e$ and $c \sqcup d = e$.
- (10) Let L be a lattice. Then the following statements are equivalent
- (i) there exists a full sublattice K of L such that M_3 and K are isomorphic,
 - (ii) there exist elements a, b, c, d, e of L such that $a \neq b$ and $a \neq c$ and $a \neq d$ and $a \neq e$ and $b \neq c$ and $b \neq d$ and $b \neq e$ and $c \neq d$ and $c \neq e$ and $d \neq e$ and $a \sqcap b = a$ and $a \sqcap c = a$ and $a \sqcap d = a$ and $b \sqcap e = b$ and $c \sqcap e = c$ and $d \sqcap e = d$ and $b \sqcap c = a$ and $b \sqcap d = a$ and $c \sqcap d = a$ and $b \sqcup c = e$ and $b \sqcup d = e$ and $c \sqcup d = e$.

3. DIAMOND AND PENTAGON

Let L be a non empty relational structure. We say that L is modular if and only if:

(Def. 3) For all elements a, b, c of L such that $a \leq c$ holds $a \sqcup (b \sqcap c) = (a \sqcup b) \sqcap c$.

One can check that every non empty antisymmetric reflexive relational structure with g.l.b.'s which is distributive is also modular.

The following two propositions are true:

- (11) Let L be a lattice. Then L is modular if and only if it is not true that there exists a full sublattice K of L such that N_5 and K are isomorphic.
- (12) Let L be a lattice. Suppose L is modular. Then L is distributive if and only if it is not true that there exists a full sublattice K of L such that M_3 and K are isomorphic.

4. INTERVALS OF A LATTICE

Let L be a non empty relational structure and let a, b be elements of L . The functor $[a, b]$ yields a subset of L and is defined by:

(Def. 4) For every element c of L holds $c \in [a, b]$ iff $a \leq c$ and $c \leq b$.

Let L be a non empty relational structure and let I_1 be a subset of L . We say that I_1 is interval if and only if:

(Def. 5) There exist elements a, b of L such that $I_1 = [a, b]$.

Let L be a non empty reflexive transitive relational structure. One can check that every subset of L which is non empty and interval is also directed and every subset of L which is non empty and interval is also filtered.

Let L be a non empty relational structure and let a, b be elements of L . One can check that $[a, b]$ is interval.

The following proposition is true

- (13) For every non empty reflexive transitive relational structure L and for all elements a, b of L holds $[a, b] = \uparrow a \cap \downarrow b$.

Let L be a poset with g.l.b.'s and let a, b be elements of L . One can check that $\text{sub}([a, b])$ is meet-inheriting.

Let L be a poset with l.u.b.'s and let a, b be elements of L . One can check that $\text{sub}([a, b])$ is join-inheriting.

We now state the proposition

- (14) Let L be a lattice and a, b be elements of L . If L is modular, then $\text{sub}([b, a \sqcup b])$ and $\text{sub}([a \sqcap b, a])$ are isomorphic.

Let us note that there exists a lattice which is finite and non empty.

Let us note that every semilattice which is finite is also lower-bounded.

Let us note that every lattice which is finite is also complete.

REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [2] Grzegorz Bancerek. Complete lattices. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/lattice3.html>.
- [3] Grzegorz Bancerek. Bounds in posets and relational substructures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_0.html.
- [4] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_0.html.
- [5] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [6] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [7] Czesław Byliński. Galois connections. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_1.html.
- [8] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.
- [9] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_1.html.
- [10] Andrzej Trybulec. Enumerated sets. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/enumset1.html>.
- [11] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [12] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/orders_1.html.
- [13] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [14] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.
- [15] Mariusz Żynel and Czesław Byliński. Properties of relational structures, posets, lattices and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_2.html.

Received April 3, 1998

Published January 2, 2004

