Ordered Rings - Part II

Michał Muzalewski Warsaw University Białystok Lesław W. Szczerba Siedlce University

Summary. This series of papers is devoted to the notion of the ordered ring, and one of its most important cases: the notion of ordered field. It follows the results of [6]. The idea of the notion of order in the ring is based on that of positive cone i.e. the set of positive elements. Positive cone has to contain at least squares of all elements, and has to be closed under sum and product. Therefore the key notions of this theory are that of square, sum of squares, product of squares, etc. and finally elements generated from squares by means of sums and products. Part II contains the classification of sums of such elements.

MML Identifier: 0_RING_2.

The terminology and notation used here are introduced in the following articles: [1], [2], [7], [3], [4], and [5]. In the sequel R is a field structure and x, y are scalars of R. One can prove the following propositions:

- (1) If x is a square and y is a square or x is a sum of squares and y is a square, then x + y is a sum of squares.
- (2) If x is a sum of products of squares and y is a square or x is a sum of products of squares and y is a product of squares, then x + y is a sum of products of squares.
- (3) If x is a amalgam of squares and y is a product of squares or x is a amalgam of squares and y is a amalgam of squares or x is a sum of amalgams of squares and y is a square or x is a sum of amalgams of squares and y is a product of squares or x is a sum of amalgams of squares and y is a amalgam of squares, then x + y is a sum of amalgams of squares.
- (4) If x is a square and y is a sum of squares or x is a square and y is a product of squares or x is a square and y is a sum of products of squares or x is a square and y is a amalgam of squares or x is a square and y is a sum of amalgams of squares or x is a square and y is generated from squares, then x + y is generated from squares.

C 1991 Fondation Philippe le Hodey ISSN 0777-4028

- (5) If x is a sum of squares and y is a sum of squares or x is a sum of squares and y is a product of squares or x is a sum of squares and y is a sum of products of squares or x is a sum of squares and y is a amalgam of squares or x is a sum of squares and y is a sum of squares or x is a sum of squares and y is a sum of squares or x is a sum of squares and y is generated from squares, then x + y is generated from squares.
- (6) If x is a product of squares and y is a square or x is a product of squares and y is a sum of squares or x is a product of squares and y is a product of squares or x is a product of squares and y is a sum of products of squares or x is a product of squares and y is a amalgam of squares or x is a product of squares and y is a sum of amalgams of squares or x is a product of squares and y is a sum of amalgams of squares or x is a product of squares and y is generated from squares, then x + y is generated from squares.
- (7) If x is a sum of products of squares and y is a sum of squares or x is a sum of products of squares and y is a sum of products of squares or x is a sum of products of squares and y is a amalgam of squares or x is a sum of products of squares and y is a sum of amalgams of squares or x is a sum of products of squares and y is generated from squares, then x + yis generated from squares.
- (8) If x is a amalgam of squares and y is a square or x is a amalgam of squares and y is a sum of squares or x is a amalgam of squares and y is a sum of products of squares or x is a amalgam of squares and y is a sum of amalgams of squares or x is a amalgam of squares and y is generated from squares, then x + y is generated from squares.
- (9) If x is a sum of amalgams of squares and y is a sum of squares or x is a sum of amalgams of squares and y is a sum of products of squares or x is a sum of amalgams of squares and y is a sum of amalgams of squares or x is a sum of amalgams of squares and y is generated from squares, then x + y is generated from squares.
- (10) If x is generated from squares and y is a square or x is generated from squares and y is a sum of squares or x is generated from squares and y is a product of squares or x is generated from squares and y is a sum of products of squares or x is generated from squares and y is a amalgam of squares or x is generated from squares and y is a sum of amalgams of squares or x is generated from squares and y is a sum of amalgams of squares or x is generated from squares and y is generated from squares, then x + y is generated from squares.

References

- Grzegorz Bancerek. The fundamental properties of natural numbers. Formalized Mathematics, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.

- Michał Muzalewski. Construction of rings and left-, right-, and bi-modules over a ring. Formalized Mathematics, 2(1):3-11, 1991.
- [5] Michał Muzalewski and Lesław W. Szczerba. Ordered rings part I. Formalized Mathematics, 2(2):243-245, 1991.
- Wanda Szmielew. From Affine to Euclidean Geometry. Volume 27, PWN D.Reidel Publ. Co., Warszawa – Dordrecht, 1983.
- [7] Andrzej Trybulec. Tarski Grothendieck set theory. Formalized Mathematics, 1(1):9–11, 1990.

Received October 11, 1990