

Several Classes of BCI-algebras and their Properties

Yuzhong Ding
Qingdao University of Science
and Technology
China

Summary. I have formalized the BCI-algebras closely following the book [6], sections 1.1 to 1.3, 1.6, 2.1 to 2.3, and 2.7. In this article the general theory of BCI-algebras and several classes of BCI-algebras are given.

MML identifier: BCIALG_1, version: 7.8.04 4.81.962

The articles [10], [4], [13], [9], [3], [12], [2], [11], [5], [7], [8], [1], and [14] provide the notation and terminology for this paper.

1. THE BASICS OF GENERAL THEORY OF BCI-ALGEBRAS

We introduce BCI structures which are extensions of 1-sorted structure and are systems

\langle a carrier, an internal complement \rangle ,

where the carrier is a set and the internal complement is a binary operation on the carrier.

Let us note that there exists a BCI structure which is non empty and strict.

Let A be a BCI structure and let x, y be elements of A . The functor $x \setminus y$ yielding an element of A is defined by:

(Def. 1) $x \setminus y = (\text{the internal complement of } A)(x, y)$.

We introduce BCI structures with 0 which are extensions of BCI structure and zero structure and are systems

\langle a carrier, an internal complement, a zero \rangle ,

where the carrier is a set, the internal complement is a binary operation on the carrier, and the zero is an element of the carrier.

Let us note that there exists a BCI structure with 0 which is non empty and strict.

Let I_1 be a non empty BCI structure with 0 and let x be an element of I_1 . The functor x^c yields an element of I_1 and is defined by:

(Def. 2) $x^c = 0_{(I_1)} \setminus x$.

Let I_1 be a non empty BCI structure with 0. We say that I_1 is B if and only if:

(Def. 3) For all elements x, y, z of I_1 holds $x \setminus y \setminus (z \setminus y) \setminus (x \setminus z) = 0_{(I_1)}$.

We say that I_1 is C if and only if:

(Def. 4) For all elements x, y, z of I_1 holds $x \setminus y \setminus z \setminus (x \setminus z \setminus y) = 0_{(I_1)}$.

We say that I_1 is I if and only if:

(Def. 5) For every element x of I_1 holds $x \setminus x = 0_{(I_1)}$.

We say that I_1 is K if and only if:

(Def. 6) For all elements x, y of I_1 holds $x \setminus y \setminus x = 0_{(I_1)}$.

We say that I_1 is BCI-4 if and only if:

(Def. 7) For all elements x, y of I_1 such that $x \setminus y = 0_{(I_1)}$ and $y \setminus x = 0_{(I_1)}$ holds $x = y$.

We say that I_1 is BCK-5 if and only if:

(Def. 8) For every element x of I_1 holds $x^c = 0_{(I_1)}$.

The BCI structure BCI-EXAMPLE with 0 is defined as follows:

(Def. 9) BCI-EXAMPLE = $\langle \{\emptyset\}, \text{op}_2, \text{op}_0 \rangle$.

Let us note that BCI-EXAMPLE is strict and non empty.

One can verify that there exists a non empty BCI structure with 0 which is strict, B, C, I, and BCI-4.

A BCI-algebra is B C I BCI-4 non empty BCI structure with 0.

Let X be a BCI-algebra. A BCI-algebra is called a subalgebra of X if it satisfies the conditions (Def. 10).

(Def. 10)(i) $0_{\text{it}} = 0_X$,

(ii) the carrier of it \subseteq the carrier of X , and

(iii) the internal complement of it = (the internal complement of X) \upharpoonright (the carrier of it).

The following proposition is true

(1) Let X be a non empty BCI structure with 0. Then X is a BCI-algebra if and only if the following conditions are satisfied:

(i) X is I and BCI-4, and

(ii) for all elements x, y, z of X holds $x \setminus y \setminus (x \setminus z) \setminus (z \setminus y) = 0_X$ and $x \setminus (x \setminus y) \setminus y = 0_X$.

One can check that there exists a BCI-algebra which is strict and BCK-5.

A BCK-algebra is BCK-5 BCI-algebra.

Let I_1 be a non empty BCI structure with 0 and let x, y be elements of I_1 .

The predicate $x \leq y$ is defined as follows:

(Def. 11) $x \setminus y = 0_{(I_1)}$.

We use the following convention: X denotes a BCI-algebra, x, y, z, u, a, b denote elements of X , and I_1 denotes a non empty subset of X .

We now state a number of propositions:

- (2) $x \setminus 0_X = x$.
- (3) If $x \setminus y = 0_X$ and $y \setminus z = 0_X$, then $x \setminus z = 0_X$.
- (4) If $x \setminus y = 0_X$, then $x \setminus z \setminus (y \setminus z) = 0_X$ and $z \setminus y \setminus (z \setminus x) = 0_X$.
- (5) If $x \leq y$, then $x \setminus z \leq y \setminus z$ and $z \setminus y \leq z \setminus x$.
- (6) If $x \setminus y = 0_X$, then $(y \setminus x)^c = 0_X$.
- (7) $x \setminus y \setminus z = x \setminus z \setminus y$.
- (8) $x \setminus (x \setminus (x \setminus y)) = x \setminus y$.
- (9) $(x \setminus y)^c = x^c \setminus y^c$.
- (10) $x \setminus (x \setminus y) \setminus (y \setminus x) \setminus (x \setminus (x \setminus (y \setminus (y \setminus x)))) = 0_X$.
- (11) Let X be a non empty BCI structure with 0. Then X is a BCI-algebra if and only if the following conditions are satisfied:
 - (i) X is BCI-4, and
 - (ii) for all elements x, y, z of X holds $x \setminus y \setminus (x \setminus z) \setminus (z \setminus y) = 0_X$ and $x \setminus 0_X = x$.
- (12) If for every BCI-algebra X and for all elements x, y of X holds $x \setminus (x \setminus y) = y \setminus (y \setminus x)$, then X is a BCK-algebra.
- (13) If for every BCI-algebra X and for all elements x, y of X holds $x \setminus y \setminus y = x \setminus y$, then X is a BCK-algebra.
- (14) If for every BCI-algebra X and for all elements x, y of X holds $x \setminus (y \setminus x) = x$, then X is a BCK-algebra.
- (15) If for every BCI-algebra X and for all elements x, y, z of X holds $(x \setminus y) \setminus y = x \setminus z \setminus (y \setminus z)$, then X is a BCK-algebra.
- (16) If for every BCI-algebra X and for all elements x, y of X holds $x \setminus y \setminus (y \setminus x) = x \setminus y$, then X is a BCK-algebra.
- (17) If for every BCI-algebra X and for all elements x, y of X holds $x \setminus y \setminus (x \setminus y \setminus (y \setminus x)) = 0_X$, then X is a BCK-algebra.
- (18) For every BCI-algebra X holds X is K iff X is a BCK-algebra.

Let X be a BCI-algebra. The functor BCK-part X yielding a non empty subset of X is defined by:

(Def. 12) BCK-part $X = \{x; x \text{ ranges over elements of } X: 0_X \leq x\}$.

Next we state the proposition

(19) $0_X \in \text{BCK-part } X$.

Let us consider X . Note that 0_X

Next we state three propositions:

(20) For all elements x, y of BCK-part X holds $x \setminus y \in \text{BCK-part } X$.

(21) For every element x of X and for every element y of BCK-part X holds $x \setminus y \leq x$.

(22) X is a subalgebra of X .

Let X be a BCI-algebra and let I_1 be a subalgebra of X . We say that I_1 is proper if and only if:

(Def. 13) $I_1 \neq X$.

Let us consider X . Note that there exists a subalgebra of X which is non proper.

Let X be a BCI-algebra and let I_1 be an element of X . We say that I_1 is atom if and only if:

(Def. 14) For every element z of X such that $z \setminus I_1 = 0_X$ holds $z = I_1$.

Let X be a BCI-algebra. The functor $\text{AtomSet } X$ yields a non empty subset of X and is defined by:

(Def. 15) $\text{AtomSet } X = \{x; x \text{ ranges over elements of } X: x \text{ is atom}\}$.

One can prove the following propositions:

(23) $0_X \in \text{AtomSet } X$.

(24) For every element x of X holds $x \in \text{AtomSet } X$ iff for every element z of X holds $z \setminus (z \setminus x) = x$.

(25) For every element x of X holds $x \in \text{AtomSet } X$ iff for all elements u, z of X holds $z \setminus u \setminus (z \setminus x) = x \setminus u$.

(26) For every element x of X holds $x \in \text{AtomSet } X$ iff for all elements y, z of X holds $x \setminus (z \setminus y) \leq y \setminus (z \setminus x)$.

(27) For every element x of X holds $x \in \text{AtomSet } X$ iff for all elements y, z, u of X holds $(x \setminus u) \setminus (z \setminus y) \leq y \setminus u \setminus (z \setminus x)$.

(28) For every element x of X holds $x \in \text{AtomSet } X$ iff for every element z of X holds $z^c \setminus x^c = x \setminus z$.

(29) For every element x of X holds $x \in \text{AtomSet } X$ iff $(x^c)^c = x$.

(30) For every element x of X holds $x \in \text{AtomSet } X$ iff for every element z of X holds $(z \setminus x)^c = x \setminus z$.

(31) For every element x of X holds $x \in \text{AtomSet } X$ iff for every element z of X holds $((x \setminus z)^c)^c = x \setminus z$.

(32) For every element x of X holds $x \in \text{AtomSet } X$ iff for all elements z, u of X holds $z \setminus (z \setminus (x \setminus u)) = x \setminus u$.

- (33) For every element a of $\text{AtomSet } X$ and for every element x of X holds $a \setminus x \in \text{AtomSet } X$.

Let X be a BCI-algebra and let a, b be elements of $\text{AtomSet } X$. Then $a \setminus b$ is an element of $\text{AtomSet } X$.

One can prove the following propositions:

- (34) For every element x of X holds $x^c \in \text{AtomSet } X$.
 (35) For every element x of X there exists an element a of $\text{AtomSet } X$ such that $a \leq x$.

Let X be a BCI-algebra. We say that X is generated by atom if and only if:

- (Def. 16) For every element x of X there exists an element a of $\text{AtomSet } X$ such that $a \leq x$.

Let X be a BCI-algebra and let a be an element of $\text{AtomSet } X$. The functor $\text{BranchV } a$ yields a non empty subset of X and is defined as follows:

- (Def. 17) $\text{BranchV } a = \{x; x \text{ ranges over elements of } X: a \leq x\}$.

We now state several propositions:

- (36) Every BCI-algebra is generated by atom.
 (37) For all elements a, b of $\text{AtomSet } X$ and for every element x of $\text{BranchV } b$ holds $a \setminus x = a \setminus b$.
 (38) For every element a of $\text{AtomSet } X$ and for every element x of $\text{BCK-part } X$ holds $a \setminus x = a$.
 (39) For all elements a, b of $\text{AtomSet } X$ and for every element x of $\text{BranchV } a$ and for every element y of $\text{BranchV } b$ holds $x \setminus y \in \text{BranchV}(a \setminus b)$.
 (40) For every element a of $\text{AtomSet } X$ and for all elements x, y of $\text{BranchV } a$ holds $x \setminus y \in \text{BCK-part } X$.
 (41) For all elements a, b of $\text{AtomSet } X$ and for every element x of $\text{BranchV } a$ and for every element y of $\text{BranchV } b$ such that $a \neq b$ holds $x \setminus y \notin \text{BCK-part } X$.
 (42) For all elements a, b of $\text{AtomSet } X$ such that $a \neq b$ holds $\text{BranchV } a \cap \text{BranchV } b = \emptyset$.

Let X be a BCI-algebra. A non empty subset of X is said to be an ideal of X if:

- (Def. 18) $0_X \in \text{it}$ and for all elements x, y of X such that $x \setminus y \in \text{it}$ and $y \in \text{it}$ holds $x \in \text{it}$.

Let X be a BCI-algebra and let I_1 be an ideal of X . We say that I_1 is closed if and only if:

- (Def. 19) For every element x of I_1 holds $x^c \in I_1$.

Let us consider X . Note that there exists an ideal of X which is closed.

Next we state four propositions:

- (43) $\{0_X\}$ is a closed ideal of X .

- (44) The carrier of X is a closed ideal of X .
- (45) BCK-part X is a closed ideal of X .
- (46) If I_1 is an ideal of X , then for all elements x, y of X such that $x \in I_1$ and $y \leq x$ holds $y \in I_1$.

2. ASSOCIATIVE BCI-ALGEBRAS

Let I_1 be a BCI-algebra. We say that I_1 is associative if and only if:

(Def. 20) For all elements x, y, z of I_1 holds $(x \setminus y) \setminus z = x \setminus (y \setminus z)$.

We say that I_1 is quasi-associative if and only if:

(Def. 21) For every element x of I_1 holds $(x^c)^c = x^c$.

We say that I_1 is positive-implicative if and only if:

(Def. 22) For all elements x, y of I_1 holds $(x \setminus (x \setminus y)) \setminus (y \setminus x) = x \setminus (x \setminus (y \setminus (y \setminus x)))$.

We say that I_1 is weakly-positive-implicative if and only if:

(Def. 23) For all elements x, y, z of I_1 holds $(x \setminus y) \setminus z = x \setminus z \setminus z \setminus (y \setminus z)$.

We say that I_1 is implicative if and only if:

(Def. 24) For all elements x, y of I_1 holds $(x \setminus (x \setminus y)) \setminus (y \setminus x) = y \setminus (y \setminus x)$.

We say that I_1 is weakly-implicative if and only if:

(Def. 25) For all elements x, y of I_1 holds $x \setminus (y \setminus x) \setminus (y \setminus x)^c = x$.

We say that I_1 is p -semisimple if and only if:

(Def. 26) For all elements x, y of I_1 holds $x \setminus (x \setminus y) = y$.

We say that I_1 is alternative if and only if:

(Def. 27) For all elements x, y of I_1 holds $x \setminus (x \setminus y) = (x \setminus x) \setminus y$ and $(x \setminus y) \setminus y = x \setminus (y \setminus y)$.

One can check that there exists a BCI-algebra which is implicative, positive-implicative, p -semisimple, associative, weakly-implicative, and weakly-positive-implicative.

Next we state several propositions:

- (47) X is associative iff for every element x of X holds $x^c = x$.
- (48) For all elements x, y of X holds $y \setminus x = x \setminus y$ iff X is associative.
- (49) Let X be a non empty BCI structure with 0. Then X is an associative BCI-algebra if and only if for all elements x, y, z of X holds $y \setminus x \setminus (z \setminus x) = z \setminus y$ and $x \setminus 0_X = x$.
- (50) Let X be a non empty BCI structure with 0. Then X is an associative BCI-algebra if and only if for all elements x, y, z of X holds $x \setminus y \setminus (x \setminus z) = z \setminus y$ and $x^c = x$.

- (51) Let X be a non empty BCI structure with 0. Then X is an associative BCI-algebra if and only if for all elements x, y, z of X holds $x \setminus y \setminus (x \setminus z) = y \setminus z$ and $x \setminus 0_X = x$.

3. p -SEMISIMPLE BCI-ALGEBRAS

One can prove the following propositions:

- (52) X is p -semisimple iff every element of X is atom.
(53) If X is p -semisimple, then BCK-part $X = \{0_X\}$.
(54) X is p -semisimple iff for every element x of X holds $(x^c)^c = x$.
(55) X is p -semisimple iff for all x, y holds $y \setminus (y \setminus x) = x$.
(56) X is p -semisimple iff for all x, y, z holds $z \setminus y \setminus (z \setminus x) = x \setminus y$.
(57) X is p -semisimple iff for all x, y, z holds $x \setminus (z \setminus y) = y \setminus (z \setminus x)$.
(58) X is p -semisimple iff for all x, y, z, u holds $(x \setminus u) \setminus (z \setminus y) = y \setminus u \setminus (z \setminus x)$.
(59) X is p -semisimple iff for all x, z holds $z^c \setminus x^c = x \setminus z$.
(60) X is p -semisimple iff for all x, z holds $((x \setminus z)^c)^c = x \setminus z$.
(61) X is p -semisimple iff for all x, u, z holds $z \setminus (z \setminus (x \setminus u)) = x \setminus u$.
(62) X is p -semisimple iff for every x such that $x^c = 0_X$ holds $x = 0_X$.
(63) X is p -semisimple iff for all x, y holds $x \setminus y^c = y \setminus x^c$.
(64) X is p -semisimple iff for all x, y, z, u holds $(x \setminus y) \setminus (z \setminus u) = x \setminus z \setminus (y \setminus u)$.
(65) X is p -semisimple iff for all x, y, z holds $x \setminus y \setminus (z \setminus y) = x \setminus z$.
(66) X is p -semisimple iff for all x, y, z holds $x \setminus (y \setminus z) = (z \setminus y) \setminus x^c$.
(67) X is p -semisimple iff for all x, y, z such that $y \setminus x = z \setminus x$ holds $y = z$.
(68) X is p -semisimple iff for all x, y, z such that $x \setminus y = x \setminus z$ holds $y = z$.
(69) Let X be a non empty BCI structure with 0. Then X is a p -semisimple BCI-algebra if and only if for all elements x, y, z of X holds $x \setminus y \setminus (x \setminus z) = z \setminus y$ and $x \setminus 0_X = x$.
(70) Let X be a non empty BCI structure with 0. Then X is a p -semisimple BCI-algebra if and only if X is I and for all elements x, y, z of X holds $x \setminus (y \setminus z) = z \setminus (y \setminus x)$ and $x \setminus 0_X = x$.

4. QUASI-ASSOCIATIVE BCI-ALGEBRAS

Next we state several propositions:

- (71) X is quasi-associative iff for every element x of X holds $x^c \leq x$.
(72) X is quasi-associative iff for all elements x, y of X holds $(x \setminus y)^c = (y \setminus x)^c$.
(73) X is quasi-associative iff for all elements x, y of X holds $x^c \setminus y = (x \setminus y)^c$.

- (74) X is quasi-associative iff for all elements x, y of X holds $x \setminus y \setminus (y \setminus x) \in$ BCK-part X .
- (75) X is quasi-associative iff for all elements x, y, z of X holds $(x \setminus y) \setminus z \leq x \setminus (y \setminus z)$.

5. ALTERNATIVE BCI-ALGEBRAS

We now state several propositions:

- (76) If X is alternative, then $x^c = x$ and $x \setminus (x \setminus y) = y$ and $x \setminus y \setminus y = x$.
- (77) If X is alternative and $x \setminus a = x \setminus b$, then $a = b$.
- (78) If X is alternative and $a \setminus x = b \setminus x$, then $a = b$.
- (79) If X is alternative and $x \setminus y = 0_X$, then $x = y$.
- (80) If X is alternative and $x \setminus a \setminus b = 0_X$, then $a = x \setminus b$ and $b = x \setminus a$.

One can check the following observations:

- * every BCI-algebra which is alternative is also associative,
- * every BCI-algebra which is associative is also alternative, and
- * every BCI-algebra which is alternative is also implicative.

The following two propositions are true:

- (81) If X is alternative, then $x \setminus (x \setminus y) \setminus (y \setminus x) = x$.
- (82) If X is alternative, then $y \setminus (y \setminus (x \setminus (x \setminus y))) = y$.

6. IMPLICATIVE, POSITIVE-IMPLICATIVE, AND WEAKLY-POSITIVE-IMPLICATIVE BCI-ALGEBRAS

Let us observe that every BCI-algebra which is associative is also weakly-positive-implicative and every BCI-algebra which is p -semisimple is also weakly-positive-implicative.

We now state two propositions:

- (83) Let X be a non empty BCI structure with 0 . Then X is an implicative BCI-algebra if and only if for all elements x, y, z of X holds $x \setminus y \setminus (x \setminus z) \setminus (z \setminus y) = 0_X$ and $x \setminus 0_X = x$ and $(x \setminus (x \setminus y)) \setminus (y \setminus x) = y \setminus (y \setminus x)$.
- (84) X is weakly-positive-implicative iff for all elements x, y of X holds $x \setminus y = x \setminus y \setminus y \setminus y^c$.

One can verify that every BCI-algebra which is positive-implicative is also weakly-positive-implicative and every BCI-algebra which is alternative is also weakly-positive-implicative.

One can prove the following two propositions:

- (85) Suppose X is a weakly-positive-implicative BCI-algebra. Let x, y be elements of X . Then $(x \setminus (x \setminus y)) \setminus (y \setminus x) = y \setminus (y \setminus x) \setminus (y \setminus x) \setminus (x \setminus y)$.

- (86) Let X be a non empty BCI structure with 0 . Then X is a positive-implicative BCI-algebra if and only if for all elements x, y, z of X holds $x \setminus y \setminus (x \setminus z) \setminus (z \setminus y) = 0_X$ and $x \setminus 0_X = x$ and $x \setminus y = x \setminus y \setminus y \setminus y^c$ and $(x \setminus (x \setminus y)) \setminus (y \setminus x) = y \setminus (y \setminus x) \setminus (y \setminus x) \setminus (x \setminus y)$.

REFERENCES

- [1] Józef Białas. Group and field definitions. *Formalized Mathematics*, 1(3):433–439, 1990.
- [2] Czesław Byliński. Binary operations. *Formalized Mathematics*, 1(1):175–180, 1990.
- [3] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [4] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [5] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Formalized Mathematics*, 1(2):335–342, 1990.
- [6] Jie Meng and YoungLin Liu. *An Introduction to BCI-algebras*. Shaanxi Scientific and Technological Press, 2001.
- [7] Michał Muzalewski. Midpoint algebras. *Formalized Mathematics*, 1(3):483–488, 1990.
- [8] Michał Muzalewski. Construction of rings and left-, right-, and bi-modules over a ring. *Formalized Mathematics*, 2(1):3–11, 1991.
- [9] Andrzej Trybulec. Subsets of complex numbers. *To appear in Formalized Mathematics*.
- [10] Andrzej Trybulec. Tarski Grothendieck set theory. *Formalized Mathematics*, 1(1):9–11, 1990.
- [11] Wojciech A. Trybulec. Groups. *Formalized Mathematics*, 1(5):821–827, 1990.
- [12] Wojciech A. Trybulec. Vectors in real linear space. *Formalized Mathematics*, 1(2):291–296, 1990.
- [13] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [14] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

Received February 23, 2007
