

# Properties of Fuzzy Relation

Noboru Endou  
Gifu National College of Technology

Takashi Mitsuishi  
Miyagi University

Keiji Ohkubo  
Shinshu University  
Nagano

**Summary.** In this article, we introduce four fuzzy relations and the composition, and some useful properties are shown by them. In section 2, the definition of converse relation  $R^{-1}$  of fuzzy relation  $R$  and properties concerning it are described. In the next section, we define the composition of the fuzzy relation and show some properties. In the final section we describe the identity relation, the universe relation and the zero relation.

MML Identifier: FUZZY\_4.

WWW: [http://mizar.org/JFM/Vol13/fuzzy\\_4.html](http://mizar.org/JFM/Vol13/fuzzy_4.html)

The articles [9], [2], [12], [1], [10], [11], [13], [5], [4], [3], [6], [7], and [8] provide the notation and terminology for this paper.

## 1. BASIC PROPERTIES OF THE MEMBERSHIP FUNCTION

We follow the rules:  $x, y, z$  denote sets and  $C_1, C_2, C_3$  denote non empty sets.

Let  $C_1$  be a non empty set and let  $F$  be a membership function of  $C_1$ . Observe that  $\text{rng } F$  is non empty.

One can prove the following four propositions:

- (1) Let  $F$  be a membership function of  $C_1$ . Then  $\text{rng } F$  is bounded and for every  $x$  such that  $x \in \text{dom } F$  holds  $F(x) \leq \sup \text{rng } F$  and for every  $x$  such that  $x \in \text{dom } F$  holds  $F(x) \geq \inf \text{rng } F$ .
- (2) For all membership functions  $F, G$  of  $C_1$  such that for every  $x$  such that  $x \in C_1$  holds  $F(x) \leq G(x)$  holds  $\sup \text{rng } F \leq \sup \text{rng } G$ .
- (3) For every membership function  $f$  of  $C_1, C_2$  and for every element  $c$  of  $[:C_1, C_2:]$  holds  $0 \leq f(c)$  and  $f(c) \leq 1$ .
- (4) For every membership function  $f$  of  $C_1, C_2$  and for all  $x, y$  such that  $\langle x, y \rangle \in [:C_1, C_2:]$  holds  $0 \leq f(\langle x, y \rangle)$  and  $f(\langle x, y \rangle) \leq 1$ .

## 2. DEFINITION OF CONVERSE FUZZY RELATION AND SOME PROPERTIES

Let  $C_1, C_2$  be non empty sets and let  $h$  be a membership function of  $C_2, C_1$ . The functor converse  $h$  yielding a membership function of  $C_1, C_2$  is defined by:

(Def. 1) For all  $x, y$  such that  $\langle x, y \rangle \in [:C_1, C_2:]$  holds  $(\text{converse } h)(\langle x, y \rangle) = h(\langle y, x \rangle)$ .

Let  $C_1, C_2$  be non empty sets, let  $f$  be a membership function of  $C_2, C_1$ , and let  $R$  be a fuzzy relation of  $C_2, C_1, f$ . The functor  $R^{-1}$  yields a fuzzy relation of  $C_1, C_2$ , converse  $f$  and is defined as follows:

(Def. 2)  $R^{-1} = [[:C_1, C_2:], (\text{converse } f)^\circ[:C_1, C_2:]]$ .

Next we state a number of propositions:

- (5) For every membership function  $f$  of  $C_1, C_2$  holds  $\text{converse } \text{converse } f = f$ .
- (6) For every membership function  $f$  of  $C_1, C_2$  and for every fuzzy relation  $R$  of  $C_1, C_2, f$  holds  $(R^{-1})^{-1} = R$ .
- (7) For every membership function  $f$  of  $C_1, C_2$  holds  $1\text{-minus } \text{converse } f = \text{converse } 1\text{-minus } f$ .
- (8) For every membership function  $f$  of  $C_1, C_2$  and for every fuzzy relation  $R$  of  $C_1, C_2, f$  holds  $(R^{-1})^c = (R^c)^{-1}$ .
- (9) For all membership functions  $f, g$  of  $C_1, C_2$  holds  $\text{converse } \max(f, g) = \max(\text{converse } f, \text{converse } g)$ .
- (10) Let  $f, g$  be membership functions of  $C_1, C_2, R$  be a fuzzy relation of  $C_1, C_2, f$ , and  $S$  be a fuzzy relation of  $C_1, C_2, g$ . Then  $(R \cup S)^{-1} = R^{-1} \cup S^{-1}$ .
- (11) For all membership functions  $f, g$  of  $C_1, C_2$  holds  $\text{converse } \min(f, g) = \min(\text{converse } f, \text{converse } g)$ .
- (12) Let  $f, g$  be membership functions of  $C_1, C_2, R$  be a fuzzy relation of  $C_1, C_2, f$ , and  $S$  be a fuzzy relation of  $C_1, C_2, g$ . Then  $(R \cap S)^{-1} = R^{-1} \cap S^{-1}$ .
- (13) Let  $f, g$  be membership functions of  $C_1, C_2$  and given  $x, y$ . If  $x \in C_1$  and  $y \in C_2$ , then if  $f(\langle x, y \rangle) \leq g(\langle x, y \rangle)$ , then  $(\text{converse } f)(\langle y, x \rangle) \leq (\text{converse } g)(\langle y, x \rangle)$ .
- (14) Let  $f, g$  be membership functions of  $C_1, C_2, R$  be a fuzzy relation of  $C_1, C_2, f$ , and  $S$  be a fuzzy relation of  $C_1, C_2, g$ . If  $R \subseteq S$ , then  $R^{-1} \subseteq S^{-1}$ .
- (15) For all membership functions  $f, g$  of  $C_1, C_2$  holds  $\text{converse } \min(f, 1\text{-minus } g) = \min(\text{converse } f, 1\text{-minus } \text{converse } g)$ .
- (16) Let  $f, g$  be membership functions of  $C_1, C_2, R$  be a fuzzy relation of  $C_1, C_2, f$ , and  $S$  be a fuzzy relation of  $C_1, C_2, g$ . Then  $(R \setminus S)^{-1} = R^{-1} \setminus S^{-1}$ .
- (17) For all membership functions  $f, g$  of  $C_1, C_2$  holds  $\text{converse } \max(\min(f, 1\text{-minus } g), \min(1\text{-minus } f, g)) = \max(\min(\text{converse } f, 1\text{-minus } \text{converse } g), \min(1\text{-minus } \text{converse } f, \text{converse } g))$ .
- (18) Let  $f, g$  be membership functions of  $C_1, C_2, R$  be a fuzzy relation of  $C_1, C_2, f$ , and  $S$  be a fuzzy relation of  $C_1, C_2, g$ . Then  $(R \dot{\setminus} S)^{-1} = R^{-1} \dot{\setminus} S^{-1}$ .

### 3. DEFINITION OF THE COMPOSITION AND SOME PROPERTIES

Let  $C_1, C_2, C_3$  be non empty sets, let  $h$  be a membership function of  $C_1, C_2$ , let  $g$  be a membership function of  $C_2, C_3$ , and let  $x, z$  be sets. Let us assume that  $x \in C_1$  and  $z \in C_3$ . The functor  $\min(h, g, x, z)$  yields a membership function of  $C_2$  and is defined by:

(Def. 3) For every element  $y$  of  $C_2$  holds  $(\min(h, g, x, z))(y) = \min(h(\langle x, y \rangle), g(\langle y, z \rangle))$ .

Let  $C_1, C_2, C_3$  be non empty sets, let  $h$  be a membership function of  $C_1, C_2$ , and let  $g$  be a membership function of  $C_2, C_3$ . The functor  $h g$  yielding a membership function of  $C_1, C_3$  is defined by:

(Def. 4) For all  $x, z$  such that  $\langle x, z \rangle \in [[:C_1, C_3:]]$  holds  $(h g)(\langle x, z \rangle) = \sup_{\text{rng}} \min(h, g, x, z)$ .

Let  $C_1, C_2, C_3$  be non empty sets, let  $f$  be a membership function of  $C_1, C_2$ , let  $g$  be a membership function of  $C_2, C_3$ , let  $R$  be a fuzzy relation of  $C_1, C_2, f$ , and let  $S$  be a fuzzy relation of  $C_2, C_3, g$ . The functor  $R S$  yielding a fuzzy relation of  $C_1, C_3, f g$  is defined by:

(Def. 5)  $RS = [:[C_1, C_3:], (fg)^\circ:[C_1, C_3:]]$ .

The following propositions are true:

- (19) For every membership function  $f$  of  $C_1, C_2$  and for all membership functions  $g, h$  of  $C_2, C_3$  holds  $f \max(g, h) = \max(fg, fh)$ .
- (20) Let  $f$  be a membership function of  $C_1, C_2, g, h$  be membership functions of  $C_2, C_3, R$  be a fuzzy relation of  $C_1, C_2, f, S$  be a fuzzy relation of  $C_2, C_3, g$ , and  $T$  be a fuzzy relation of  $C_2, C_3, h$ . Then  $R(S \cup T) = RS \cup RT$ .
- (21) For all membership functions  $f, g$  of  $C_1, C_2$  and for every membership function  $h$  of  $C_2, C_3$  holds  $\max(f, g)h = \max(fh, gh)$ .
- (22) Let  $f, g$  be membership functions of  $C_1, C_2, h$  be a membership function of  $C_2, C_3, R$  be a fuzzy relation of  $C_1, C_2, f, S$  be a fuzzy relation of  $C_1, C_2, g$ , and  $T$  be a fuzzy relation of  $C_2, C_3, h$ . Then  $(R \cup S)T = RT \cup ST$ .
- (23) Let  $f$  be a membership function of  $C_1, C_2, g, h$  be membership functions of  $C_2, C_3$ , and  $x, z$  be sets. If  $x \in C_1$  and  $z \in C_3$ , then  $(f \min(g, h))(\langle x, z \rangle) \leq (\min(fg, fh))(\langle x, z \rangle)$ .
- (24) Let  $f$  be a membership function of  $C_1, C_2, g, h$  be membership functions of  $C_2, C_3, R$  be a fuzzy relation of  $C_1, C_2, f, S$  be a fuzzy relation of  $C_2, C_3, g$ , and  $T$  be a fuzzy relation of  $C_2, C_3, h$ . Then  $R(S \cap T) \subseteq (RS) \cap (RT)$ .
- (25) Let  $f, g$  be membership functions of  $C_1, C_2, h$  be a membership function of  $C_2, C_3$ , and  $x, z$  be sets. If  $x \in C_1$  and  $z \in C_3$ , then  $(\min(f, g)h)(\langle x, z \rangle) \leq (\min(fh, gh))(\langle x, z \rangle)$ .
- (26) Let  $f, g$  be membership functions of  $C_1, C_2, h$  be a membership function of  $C_2, C_3, R$  be a fuzzy relation of  $C_1, C_2, f, S$  be a fuzzy relation of  $C_1, C_2, g$ , and  $T$  be a fuzzy relation of  $C_2, C_3, h$ . Then  $(R \cap S)T \subseteq (RT) \cap (ST)$ .
- (27) For every membership function  $f$  of  $C_1, C_2$  and for every membership function  $g$  of  $C_2, C_3$  holds converse  $fg = \text{converse } g \text{ converse } f$ .
- (28) Let  $f$  be a membership function of  $C_1, C_2, g$  be a membership function of  $C_2, C_3, R$  be a fuzzy relation of  $C_1, C_2, f$ , and  $S$  be a fuzzy relation of  $C_2, C_3, g$ . Then  $(RS)^{-1} = S^{-1}R^{-1}$ .
- (29) Let  $f, g$  be membership functions of  $C_1, C_2, h, k$  be membership functions of  $C_2, C_3$ , and  $x, z$  be sets. Suppose  $x \in C_1$  and  $z \in C_3$  and for every set  $y$  such that  $y \in C_2$  holds  $f(\langle x, y \rangle) \leq g(\langle x, y \rangle)$  and  $h(\langle y, z \rangle) \leq k(\langle y, z \rangle)$ . Then  $(fh)(\langle x, z \rangle) \leq (gk)(\langle x, z \rangle)$ .
- (30) Let  $f, g$  be membership functions of  $C_1, C_2, h, k$  be membership functions of  $C_2, C_3, R$  be a fuzzy relation of  $C_1, C_2, f, S$  be a fuzzy relation of  $C_1, C_2, g, T$  be a fuzzy relation of  $C_2, C_3, h$ , and  $W$  be a fuzzy relation of  $C_2, C_3, k$ . If  $R \subseteq S$  and  $T \subseteq W$ , then  $RT \subseteq SW$ .

#### 4. DEFINITION OF IDENTITY RELATION AND PROPERTIES OF UNIVERSE AND ZERO RELATION

Let  $C_1, C_2$  be non empty sets. The functor  $\text{Imf}(C_1, C_2)$  yielding a membership function of  $C_1, C_2$  is defined as follows:

(Def. 6) For all  $x, y$  such that  $\langle x, y \rangle \in [C_1, C_2:]$  holds if  $x = y$ , then  $(\text{Imf}(C_1, C_2))(\langle x, y \rangle) = 1$  and if  $x \neq y$ , then  $(\text{Imf}(C_1, C_2))(\langle x, y \rangle) = 0$ .

Next we state several propositions:

- (31) For every element  $c$  of  $[C_1, C_2:]$  holds  $(\text{Zmf}(C_1, C_2))(c) = 0$  and  $(\text{Umf}(C_1, C_2))(c) = 1$ .
- (32) For all  $x, y$  such that  $\langle x, y \rangle \in [C_1, C_2:]$  holds  $(\text{Zmf}(C_1, C_2))(\langle x, y \rangle) = 0$  and  $(\text{Umf}(C_1, C_2))(\langle x, y \rangle) = 1$ .

- (33) Let  $f$  be a membership function of  $C_2, C_3$ ,  $O_1$  be a zero relation of  $C_1, C_2$ ,  $O_2$  be a zero relation of  $C_1, C_3$ , and  $R$  be a fuzzy relation of  $C_2, C_3, f$ . Then  $O_1 R = O_2$ .
- (34) For every membership function  $f$  of  $C_1, C_2$  holds  $f \text{ Zmf}(C_2, C_3) = \text{Zmf}(C_1, C_3)$ .
- (35) Let  $f$  be a membership function of  $C_1, C_2$ ,  $O_1$  be a zero relation of  $C_2, C_3$ ,  $O_2$  be a zero relation of  $C_1, C_3$ , and  $R$  be a fuzzy relation of  $C_1, C_2, f$ . Then  $R O_1 = O_2$ .
- (36) For every membership function  $f$  of  $C_1, C_1$  holds  $f \text{ Zmf}(C_1, C_1) = \text{Zmf}(C_1, C_1) f$ .
- (37) Let  $f$  be a membership function of  $C_1, C_1$ ,  $O$  be a zero relation of  $C_1, C_1$ , and  $R$  be a fuzzy relation of  $C_1, C_1, f$ . Then  $R O = O R$ .

## REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Voll/ordinal1.html>.
- [2] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/zfmisc\\_1.html](http://mizar.org/JFM/Voll/zfmisc_1.html).
- [3] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in  $E^2$ . *Journal of Formalized Mathematics*, 9, 1997. [http://mizar.org/JFM/Vol9/pscomp\\_1.html](http://mizar.org/JFM/Vol9/pscomp_1.html).
- [4] Jarosław Kotowicz. Convergent real sequences. Upper and lower bound of sets of real numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/seq\\_4.html](http://mizar.org/JFM/Voll/seq_4.html).
- [5] Jarosław Kotowicz. Real sequences and basic operations on them. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/seq\\_1.html](http://mizar.org/JFM/Voll/seq_1.html).
- [6] Takashi Mitsuishi, Noboru Endou, and Yasunari Shidama. The concept of fuzzy set and membership function and basic properties of fuzzy set operation. *Journal of Formalized Mathematics*, 12, 2000. [http://mizar.org/JFM/Voll12/fuzzy\\_1.html](http://mizar.org/JFM/Voll12/fuzzy_1.html).
- [7] Takashi Mitsuishi, Katsumi Wasaki, and Yasunari Shidama. Basic properties of fuzzy set operation and membership function. *Journal of Formalized Mathematics*, 12, 2000. [http://mizar.org/JFM/Voll12/fuzzy\\_2.html](http://mizar.org/JFM/Voll12/fuzzy_2.html).
- [8] Takashi Mitsuishi, Katsumi Wasaki, and Yasunari Shidama. The concept of fuzzy relation and basic properties of its operation. *Journal of Formalized Mathematics*, 12, 2000. [http://mizar.org/JFM/Voll12/fuzzy\\_3.html](http://mizar.org/JFM/Voll12/fuzzy_3.html).
- [9] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [10] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [11] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers operations: min, max, square, and square root. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/square\\_1.html](http://mizar.org/JFM/Voll/square_1.html).
- [12] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/subset\\_1.html](http://mizar.org/JFM/Voll/subset_1.html).
- [13] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/relset\\_1.html](http://mizar.org/JFM/Voll/relset_1.html).

Received June 25, 2001

Published January 2, 2004

---