

# On the Decomposition of a Simple Closed Curve into Two Arcs

Andrzej Trybulec  
University of Białystok

Yatsuka Nakamura  
Shinshu University  
Nagano

**Summary.** The purpose of the paper is to prove lemmas needed for the Jordan curve theorem. The main result is that the decomposition of a simple closed curve into two arcs with the ends  $p_1, p_2$  is unique in the sense that every arc on the curve with the same ends must be equal to one of them.

MML Identifier: JORDAN16.

WWW: <http://mizar.org/JFM/Vol14/jordan16.html>

The articles [25], [24], [27], [1], [26], [28], [3], [5], [9], [4], [22], [17], [21], [8], [7], [20], [2], [23], [15], [10], [6], [11], [19], [18], [12], [14], [13], and [16] provide the notation and terminology for this paper.

One can prove the following proposition

- (1) Let  $S_1$  be a finite non empty subset of  $\mathbb{R}$  and  $e$  be a real number. If for every real number  $r$  such that  $r \in S_1$  holds  $r < e$ , then  $\max S_1 < e$ .

For simplicity, we follow the rules:  $C$  is a simple closed curve,  $A, A_1, A_2$  are subsets of  $\mathcal{E}_T^2$ ,  $p, p_1, p_2, q, q_1, q_2$  are points of  $\mathcal{E}_T^2$ , and  $n$  is a natural number.

Let us consider  $n$ . Observe that there exists a subset of  $\mathcal{E}_T^n$  which is trivial.

One can prove the following propositions:

- (2) For all sets  $a, b, c, X$  such that  $a \in X$  and  $b \in X$  and  $c \in X$  holds  $\{a, b, c\} \subseteq X$ .
- (3)  $\emptyset_{\mathcal{E}_T^n}$  is Bounded.
- (4)  $\text{LowerArc}(C) \neq \text{UpperArc}(C)$ .
- (5)  $\text{Segment}(A, p_1, p_2, q_1, q_2) \subseteq A$ .
- (6) For every non empty topological space  $T$  and for all subsets  $A, B$  of  $T$  such that  $A \subseteq B$  holds  $T \setminus A$  is a subspace of  $T \setminus B$ .
- (7) If  $A$  is an arc from  $p_1$  to  $p_2$  and  $q \in A$ , then  $q \in \text{LSegment}(A, p_1, p_2, q)$ .
- (8) If  $A$  is an arc from  $p_1$  to  $p_2$  and  $q \in A$ , then  $q \in \text{RSegment}(A, p_1, p_2, q)$ .
- (9) If  $A$  is an arc from  $p_1$  to  $p_2$  and  $\text{LE } q_1, q_2, A, p_1, p_2$ , then  $q_1 \in \text{Segment}(A, p_1, p_2, q_1, q_2)$  and  $q_2 \in \text{Segment}(A, p_1, p_2, q_1, q_2)$ .
- (10)  $\text{Segment}(p, q, C) \subseteq C$ .

- (11) If  $p \in C$  and  $q \in C$ , then  $p \leq_C q$  or  $q \leq_C p$ .
- (12) Let  $X, Y$  be non empty topological spaces,  $Y_0$  be a non empty subspace of  $Y$ ,  $f$  be a map from  $X$  into  $Y$ , and  $g$  be a map from  $X$  into  $Y_0$ . If  $f = g$  and  $f$  is continuous, then  $g$  is continuous.
- (13) Let  $S, T$  be non empty topological spaces,  $S_0$  be a non empty subspace of  $S$ ,  $T_0$  be a non empty subspace of  $T$ , and  $f$  be a map from  $S$  into  $T$ . Suppose  $f$  is a homeomorphism. Let  $g$  be a map from  $S_0$  into  $T_0$ . If  $g = f|_{S_0}$  and  $g$  is onto, then  $g$  is a homeomorphism.
- (14) Let  $P_1, P_2, P_3$  be subsets of  $\mathcal{E}_T^2$  and  $p_1, p_2$  be points of  $\mathcal{E}_T^2$ . Suppose  $P_1$  is an arc from  $p_1$  to  $p_2$  and  $P_2$  is an arc from  $p_1$  to  $p_2$  and  $P_3$  is an arc from  $p_1$  to  $p_2$  and  $P_2 \cap P_3 = \{p_1, p_2\}$  and  $P_1 \subseteq P_2 \cup P_3$ . Then  $P_1 = P_2$  or  $P_1 = P_3$ .
- (15) Let  $C$  be a simple closed curve,  $A_1, A_2$  be subsets of  $\mathcal{E}_T^2$ , and  $p_1, p_2$  be points of  $\mathcal{E}_T^2$ . Suppose  $A_1$  is an arc from  $p_1$  to  $p_2$  and  $A_2$  is an arc from  $p_1$  to  $p_2$  and  $A_1 \subseteq C$  and  $A_2 \subseteq C$  and  $A_1 \neq A_2$ . Then  $A_1 \cup A_2 = C$  and  $A_1 \cap A_2 = \{p_1, p_2\}$ .
- (16) Let  $A_1, A_2$  be subsets of  $\mathcal{E}_T^2$  and  $p_1, p_2, q_1, q_2$  be points of  $\mathcal{E}_T^2$ . If  $A_1$  is an arc from  $p_1$  to  $p_2$  and  $A_1 \cap A_2 = \{q_1, q_2\}$ , then  $A_1 \neq A_2$ .
- (17) Let  $C$  be a simple closed curve,  $A_1, A_2$  be subsets of  $\mathcal{E}_T^2$ , and  $p_1, p_2$  be points of  $\mathcal{E}_T^2$ . Suppose  $A_1$  is an arc from  $p_1$  to  $p_2$  and  $A_2$  is an arc from  $p_1$  to  $p_2$  and  $A_1 \subseteq C$  and  $A_2 \subseteq C$  and  $A_1 \cap A_2 = \{p_1, p_2\}$ . Then  $A_1 \cup A_2 = C$ .
- (18) Suppose  $A_1 \subseteq C$  and  $A_2 \subseteq C$  and  $A_1 \neq A_2$  and  $A_1$  is an arc from  $p_1$  to  $p_2$  and  $A_2$  is an arc from  $p_1$  to  $p_2$ . Let given  $A$ . If  $A$  is an arc from  $p_1$  to  $p_2$  and  $A \subseteq C$ , then  $A = A_1$  or  $A = A_2$ .
- (19) Let  $C$  be a simple closed curve and  $A$  be a non empty subset of  $\mathcal{E}_T^2$ . If  $A$  is an arc from  $W_{\min}(C)$  to  $E_{\max}(C)$  and  $A \subseteq C$ , then  $A = \text{LowerArc}(C)$  or  $A = \text{UpperArc}(C)$ .
- (20) Suppose  $A$  is an arc from  $p_1$  to  $p_2$  and LE  $q_1, q_2, A, p_1, p_2$ . Then there exists a map  $g$  from  $\mathbb{I}$  into  $(\mathcal{E}_T^2)|_A$  and there exist real numbers  $s_1, s_2$  such that  $g$  is a homeomorphism and  $g(0) = p_1$  and  $g(1) = p_2$  and  $g(s_1) = q_1$  and  $g(s_2) = q_2$  and  $0 \leq s_1$  and  $s_1 \leq s_2$  and  $s_2 \leq 1$ .
- (21) Suppose  $A$  is an arc from  $p_1$  to  $p_2$  and LE  $q_1, q_2, A, p_1, p_2$  and  $q_1 \neq q_2$ . Then there exists a map  $g$  from  $\mathbb{I}$  into  $(\mathcal{E}_T^2)|_A$  and there exist real numbers  $s_1, s_2$  such that  $g$  is a homeomorphism and  $g(0) = p_1$  and  $g(1) = p_2$  and  $g(s_1) = q_1$  and  $g(s_2) = q_2$  and  $0 \leq s_1$  and  $s_1 < s_2$  and  $s_2 \leq 1$ .
- (22) If  $A$  is an arc from  $p_1$  to  $p_2$  and LE  $q_1, q_2, A, p_1, p_2$ , then  $\text{Segment}(A, p_1, p_2, q_1, q_2)$  is non empty.
- (23) If  $p \in C$ , then  $p \in \text{Segment}(p, W_{\min}(C), C)$  and  $W_{\min}(C) \in \text{Segment}(p, W_{\min}(C), C)$ .

Let  $f$  be a partial function from  $\mathbb{R}$  to  $\mathbb{R}$ . We say that  $f$  is continuous if and only if:

(Def. 1)  $f$  is continuous on  $\text{dom } f$ .

Let  $f$  be a function from  $\mathbb{R}$  into  $\mathbb{R}$ . Let us observe that  $f$  is continuous if and only if:

(Def. 2)  $f$  is continuous on  $\mathbb{R}$ .

Let  $a, b$  be real numbers. The functor  $\text{AffineMap}(a, b)$  yielding a function from  $\mathbb{R}$  into  $\mathbb{R}$  is defined by:

(Def. 3) For every real number  $x$  holds  $(\text{AffineMap}(a, b))(x) = a \cdot x + b$ .

Let  $a, b$  be real numbers. One can check that  $\text{AffineMap}(a, b)$  is continuous.

Let us observe that there exists a function from  $\mathbb{R}$  into  $\mathbb{R}$  which is continuous.

Next we state a number of propositions:

- (24) Let  $f, g$  be continuous partial functions from  $\mathbb{R}$  to  $\mathbb{R}$ . Then  $g \cdot f$  is a continuous partial function from  $\mathbb{R}$  to  $\mathbb{R}$ .

- (25) For all real numbers  $a, b$  holds  $(\text{AffineMap}(a, b))(0) = b$ .
- (26) For all real numbers  $a, b$  holds  $(\text{AffineMap}(a, b))(1) = a + b$ .
- (27) For all real numbers  $a, b$  such that  $a \neq 0$  holds  $\text{AffineMap}(a, b)$  is one-to-one.
- (28) For all real numbers  $a, b, x, y$  such that  $a > 0$  and  $x < y$  holds  $(\text{AffineMap}(a, b))(x) < (\text{AffineMap}(a, b))(y)$ .
- (29) For all real numbers  $a, b, x, y$  such that  $a < 0$  and  $x < y$  holds  $(\text{AffineMap}(a, b))(x) > (\text{AffineMap}(a, b))(y)$ .
- (30) For all real numbers  $a, b, x, y$  such that  $a \geq 0$  and  $x \leq y$  holds  $(\text{AffineMap}(a, b))(x) \leq (\text{AffineMap}(a, b))(y)$ .
- (31) For all real numbers  $a, b, x, y$  such that  $a \leq 0$  and  $x \leq y$  holds  $(\text{AffineMap}(a, b))(x) \geq (\text{AffineMap}(a, b))(y)$ .
- (32) For all real numbers  $a, b$  such that  $a \neq 0$  holds  $\text{rng AffineMap}(a, b) = \mathbb{R}$ .
- (33) For all real numbers  $a, b$  such that  $a \neq 0$  holds  $(\text{AffineMap}(a, b))^{-1} = \text{AffineMap}(a^{-1}, -\frac{b}{a})$ .
- (34) For all real numbers  $a, b$  such that  $a > 0$  holds  $(\text{AffineMap}(a, b))^{\circ}[0, 1] = [b, a + b]$ .
- (35) For every map  $f$  from  $\mathbb{R}^1$  into  $\mathbb{R}^1$  and for all real numbers  $a, b$  such that  $a \neq 0$  and  $f = \text{AffineMap}(a, b)$  holds  $f$  is a homeomorphism.
- (36) If  $A$  is an arc from  $p_1$  to  $p_2$  and LE  $q_1, q_2, A, p_1, p_2$  and  $q_1 \neq q_2$ , then  $\text{Segment}(A, p_1, p_2, q_1, q_2)$  is an arc from  $q_1$  to  $q_2$ .
- (37) Let  $p_1, p_2$  be points of  $\mathcal{E}_T^2$  and  $P$  be a subset of  $\mathcal{E}_T^2$ . Suppose  $P \subseteq C$  and  $P$  is an arc from  $p_1$  to  $p_2$  and  $W_{\min}(C) \in P$  and  $E_{\max}(C) \in P$ . Then  $\text{UpperArc}(C) \subseteq P$  or  $\text{LowerArc}(C) \subseteq P$ .

## REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [2] Leszek Borys. Paracompact and metrizable spaces. *Journal of Formalized Mathematics*, 3, 1991. [http://mizar.org/JFM/Vol3/pcomps\\_1.html](http://mizar.org/JFM/Vol3/pcomps_1.html).
- [3] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_1.html](http://mizar.org/JFM/Vol1/funct_1.html).
- [4] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_2.html](http://mizar.org/JFM/Vol1/funct_2.html).
- [5] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/partfun1.html>.
- [6] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in  $\mathcal{E}^2$ . *Journal of Formalized Mathematics*, 9, 1997. [http://mizar.org/JFM/Vol9/pscomp\\_1.html](http://mizar.org/JFM/Vol9/pscomp_1.html).
- [7] Agata Darmochwał. Compact spaces. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/compts\\_1.html](http://mizar.org/JFM/Vol1/compts_1.html).
- [8] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/tops\\_2.html](http://mizar.org/JFM/Vol1/tops_2.html).
- [9] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finset\\_1.html](http://mizar.org/JFM/Vol1/finset_1.html).
- [10] Agata Darmochwał. The Euclidean space. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/euclid.html>.
- [11] Agata Darmochwał and Yatsuka Nakamura. Metric spaces as topological spaces — fundamental concepts. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/topmetr.html>.
- [12] Agata Darmochwał and Yatsuka Nakamura. The topological space  $\mathcal{E}_T^2$ . Arcs, line segments and special polygonal arcs. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/topreal1.html>.
- [13] Agata Darmochwał and Yatsuka Nakamura. The topological space  $\mathcal{E}_T^2$ . Simple closed curves. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/topreal2.html>.
- [14] Adam Grabowski and Yatsuka Nakamura. The ordering of points on a curve. Part II. *Journal of Formalized Mathematics*, 9, 1997. <http://mizar.org/JFM/Vol9/jordan5c.html>.

- [15] Zbigniew Karno. Continuity of mappings over the union of subspaces. *Journal of Formalized Mathematics*, 4, 1992. [http://mizar.org/JFM/Vol4/tmap\\_1.html](http://mizar.org/JFM/Vol4/tmap_1.html).
- [16] Yatsuka Nakamura. On the dividing function of the simple closed curve into segments. *Journal of Formalized Mathematics*, 10, 1998. <http://mizar.org/JFM/Vol10/jordan7.html>.
- [17] Yatsuka Nakamura, Piotr Rudnicki, Andrzej Trybulec, and Pauline N. Kawamoto. Preliminaries to circuits, I. *Journal of Formalized Mathematics*, 6, 1994. [http://mizar.org/JFM/Vol6/pre\\_circ.html](http://mizar.org/JFM/Vol6/pre_circ.html).
- [18] Yatsuka Nakamura and Andrzej Trybulec. A decomposition of simple closed curves and the order of their points. *Journal of Formalized Mathematics*, 9, 1997. <http://mizar.org/JFM/Vol9/jordan6.html>.
- [19] Yatsuka Nakamura, Andrzej Trybulec, and Czesław Byliński. Bounded domains and unbounded domains. *Journal of Formalized Mathematics*, 11, 1999. <http://mizar.org/JFM/Vol11/jordan2c.html>.
- [20] Beata Padlewska. Locally connected spaces. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/connsp\\_2.html](http://mizar.org/JFM/Vol2/connsp_2.html).
- [21] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/pre\\_topc.html](http://mizar.org/JFM/Vol1/pre_topc.html).
- [22] Konrad Raczkowski and Paweł Sadowski. Real function continuity. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/fcont\\_1.html](http://mizar.org/JFM/Vol2/fcont_1.html).
- [23] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/rcomp\\_1.html](http://mizar.org/JFM/Vol2/rcomp_1.html).
- [24] Andrzej Trybulec. Enumerated sets. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/enumset1.html>.
- [25] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [26] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [27] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/subset\\_1.html](http://mizar.org/JFM/Vol1/subset_1.html).
- [28] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relat\\_1.html](http://mizar.org/JFM/Vol1/relat_1.html).

*Received September 16, 2002*

*Published January 2, 2004*

---