

Some Properties of Dyadic Numbers and Intervals

Józef Białas
Łódź University

Yatsuka Nakamura
Shinshu University
Nagano

Summary. The article is the second part of a paper proving the fundamental Urysohn Theorem concerning the existence of a real valued continuous function on a normal topological space. The paper is divided into two parts. In the first part, we introduce some definitions and theorems concerning properties of intervals; in the second we prove some of properties of dyadic numbers used in proving Urysohn Lemma.

MML Identifier: URYSOHN2.

WWW: <http://mizar.org/JFM/Vol13/urysohn2.html>

The articles [11], [13], [1], [8], [12], [9], [2], [3], [4], [5], [6], [10], and [7] provide the notation and terminology for this paper.

One can prove the following propositions:

- (1) For every interval A such that $A \neq \emptyset$ holds if $\inf A < \sup A$, then $\text{vol}(A) = \sup A - \inf A$ and if $\sup A = \inf A$, then $\text{vol}(A) = 0_{\mathbb{R}}$.
- (2) For every subset A of \mathbb{R} and for every real number x such that $x \neq 0$ holds $x^{-1} \cdot (x \cdot A) = A$.
- (3) For every real number x such that $x \neq 0$ and for every subset A of \mathbb{R} such that $A = \mathbb{R}$ holds $x \cdot A = A$.
- (4) For every subset A of \mathbb{R} such that $A \neq \emptyset$ holds $0 \cdot A = \{0\}$.
- (5) For every real number x holds $x \cdot 0 = 0$.
- (6) Let a, b be extended real numbers. Suppose $a \leq b$. Then $a = -\infty$ and $b = -\infty$ or $a = -\infty$ and $b \in \mathbb{R}$ or $a = -\infty$ and $b = +\infty$ or $a \in \mathbb{R}$ and $b \in \mathbb{R}$ or $a \in \mathbb{R}$ and $b = +\infty$ or $a = +\infty$ and $b = +\infty$.
- (7) For every extended real number x holds $[x, x]$ is an interval.
- (8) For every interval A holds $0 \cdot A$ is an interval.
- (9) Let A be an interval and x be a real number. If $x \neq 0$, then if A is open interval, then $x \cdot A$ is open interval.
- (10) Let A be an interval and x be a real number. If $x \neq 0$, then if A is closed interval, then $x \cdot A$ is closed interval.
- (11) Let A be an interval and x be a real number. Suppose $0 < x$. If A is right open interval, then $x \cdot A$ is right open interval.
- (12) Let A be an interval and x be a real number. Suppose $x < 0$. If A is right open interval, then $x \cdot A$ is left open interval.

- (13) Let A be an interval and x be a real number. Suppose $0 < x$. If A is left open interval, then $x \cdot A$ is left open interval.
- (14) Let A be an interval and x be a real number. Suppose $x < 0$. If A is left open interval, then $x \cdot A$ is right open interval.
- (15) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A = [\inf A, \sup A]$. Then $B = [\inf B, \sup B]$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.
- (16) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A =]\inf A, \sup A]$. Then $B =]\inf B, \sup B]$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.
- (17) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A =]\inf A, \sup A[$. Then $B =]\inf B, \sup B[$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.
- (18) Let A be an interval. Suppose $A \neq \emptyset$. Let x be a real number. Suppose $0 < x$. Let B be an interval. Suppose $B = x \cdot A$. Suppose $A = [\inf A, \sup A[$. Then $B = [\inf B, \sup B[$ and for all real numbers s, t such that $s = \inf A$ and $t = \sup A$ holds $\inf B = x \cdot s$ and $\sup B = x \cdot t$.
- (19) For every interval A and for every real number x holds $x \cdot A$ is an interval.

Let A be an interval and let x be a real number. Note that $x \cdot A$ is interval.

One can prove the following propositions:

- (20) Let A be an interval and x be a real number. If $0 \leq x$, then for every real number y such that $y = \text{vol}(A)$ holds $x \cdot y = \text{vol}(x \cdot A)$.
- (23)¹ For every real number e_1 such that $0 < e_1$ there exists a natural number n such that $1 < 2^n \cdot e_1$.
- (24) For all real numbers a, b such that $0 \leq a$ and $1 < b - a$ there exists a natural number n such that $a < n$ and $n < b$.
- (27)² For every natural number n holds $\text{dyadic}(n) \subseteq \text{DYADIC}$.
- (28) For all real numbers a, b such that $a < b$ and $0 \leq a$ and $b \leq 1$ there exists a real number c such that $c \in \text{DYADIC}$ and $a < c$ and $c < b$.
- (29) For all real numbers a, b such that $a < b$ there exists a real number c such that $c \in \text{DOM}$ and $a < c$ and $c < b$.
- (30) For every non empty subset A of $\overline{\mathbb{R}}$ and for all extended real numbers a, b such that $A \subseteq [a, b]$ holds $a \leq \inf A$ and $\sup A \leq b$.
- (31) $0 \in \text{DYADIC}$ and $1 \in \text{DYADIC}$.
- (32) For all extended real numbers a, b such that $a = 0$ and $b = 1$ holds $\text{DYADIC} \subseteq [a, b]$.
- (33) For all natural numbers n, k such that $n \leq k$ holds $\text{dyadic}(n) \subseteq \text{dyadic}(k)$.
- (34) For all real numbers a, b, c, d such that $a < c$ and $c < b$ and $a < d$ and $d < b$ holds $|d - c| < b - a$.
- (35) Let e_1 be a real number. Suppose $0 < e_1$. Let d be a real number. Suppose $0 < d$ and $d \leq 1$. Then there exist real numbers r_1, r_2 such that $r_1 \in \text{DYADIC} \cup \mathbb{R}_{>1}$ and $r_2 \in \text{DYADIC} \cup \overline{\mathbb{R}}_{>1}$ and $0 < r_1$ and $r_1 < d$ and $d < r_2$ and $r_2 - r_1 < e_1$.

¹ The propositions (21) and (22) have been removed.

² The propositions (25) and (26) have been removed.

REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [2] Józef Białas. Infimum and supremum of the set of real numbers. Measure theory. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/supinf_1.html.
- [3] Józef Białas. Series of positive real numbers. Measure theory. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/supinf_2.html.
- [4] Józef Białas. Properties of the intervals of real numbers. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/measure5.html>.
- [5] Józef Białas. Some properties of the intervals. *Journal of Formalized Mathematics*, 6, 1994. <http://mizar.org/JFM/Vol6/measure6.html>.
- [6] Józef Białas and Yatsuka Nakamura. Dyadic numbers and T_4 topological spaces. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/urysohn1.html>.
- [7] Noboru Endou, Katsumi Wasaki, and Yasunari Shidama. Scalar multiple of Riemann definite integral. *Journal of Formalized Mathematics*, 11, 1999. <http://mizar.org/JFM/Vol11/integra2.html>.
- [8] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/real_1.html.
- [9] Rafał Kwiatek. Factorial and Newton coefficients. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/newton.html>.
- [10] Jan Popiołek. Some properties of functions modul and signum. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/absvalue.html>.
- [11] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [12] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [13] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

Received February 16, 2001

Published January 2, 2004
