

Representation Theorem for Free Continuous Lattices

Piotr Rudnicki
University of Alberta
Edmonton

Summary. We present the Mizar formalization of theorem 4.17, Chapter I from [13]: a free continuous lattice with m generators is isomorphic to the lattice of filters of 2^X ($\overline{X} = m$) which is freely generated by $\{\uparrow x : x \in X\}$ (the set of ultrafilters).

MML Identifier: WAYBEL22.

WWW: <http://mizar.org/JFM/Vol10/waybel22.html>

The articles [20], [10], [25], [18], [26], [8], [9], [3], [12], [16], [1], [2], [19], [24], [4], [22], [23], [17], [21], [5], [14], [27], [6], [11], [7], and [15] provide the notation and terminology for this paper.

1. PRELIMINARIES

One can prove the following propositions:

- (1) For every upper-bounded semilattice L and for every non empty directed subset F of $\langle \text{Filt}(L), \subseteq \rangle$ holds $\sup F = \bigcup F$.
- (2) Let L, S, T be complete non empty posets, f be a CLHomomorphism of L, S , and g be a CLHomomorphism of S, T . Then $g \cdot f$ is a CLHomomorphism of L, T .
- (3) For every non empty relational structure L holds id_L is infs-preserving.
- (4) For every non empty relational structure L holds id_L is directed-sups-preserving.
- (5) For every complete non empty poset L holds id_L is a CLHomomorphism of L, L .
- (6) For every upper-bounded non empty poset L with g.l.b.'s holds $\langle \text{Filt}(L), \subseteq \rangle$ is a continuous subframe of $2_{\subseteq}^{\text{the carrier of } L}$.

Let L be an upper-bounded non empty poset with g.l.b.'s. Observe that $\langle \text{Filt}(L), \subseteq \rangle$ is continuous.

Let L be an upper-bounded non empty poset. One can check that every element of $\langle \text{Filt}(L), \subseteq \rangle$ is non empty.

2. FREE GENERATORS OF CONTINUOUS LATTICES

Let S be a continuous complete non empty poset and let A be a set. We say that A is a set of free generators of S if and only if the condition (Def. 1) is satisfied.

(Def. 1) Let T be a continuous complete non empty poset and f be a function from A into the carrier of T . Then there exists a CLHomomorphism h of S, T such that $h \upharpoonright A = f$ and for every CLHomomorphism h' of S, T such that $h' \upharpoonright A = f$ holds $h' = h$.

The following propositions are true:

(7) Let S be a continuous complete non empty poset and A be a set. If A is a set of free generators of S , then A is a subset of S .

(8) Let S be a continuous complete non empty poset and A be a set. Suppose A is a set of free generators of S . Let h' be a CLHomomorphism of S, S . If $h' \upharpoonright A = \text{id}_A$, then $h' = \text{id}_S$.

3. REPRESENTATION THEOREM FOR FREE CONTINUOUS LATTICES

In the sequel X denotes a set, F denotes a filter of 2_{\subseteq}^X , x denotes an element of 2_{\subseteq}^X , and z denotes an element of X .

Let us consider X . The fixed ultrafilters of X is a family of subsets of 2_{\subseteq}^X and is defined as follows:

(Def. 2) The fixed ultrafilters of $X = \{\uparrow x : \bigvee_z x = \{z\}\}$.

Next we state three propositions:

(9) The fixed ultrafilters of $X \subseteq \text{Filt}(2_{\subseteq}^X)$.

(10) $\overline{\overline{\text{the fixed ultrafilters of } X}} = \overline{X}$.

(11) $F = \bigsqcup_{(Y \in F)} \{ \bigcap_{(x \in Filt(2_{\subseteq}^X), \subseteq)} \{ \uparrow x : \bigvee_z (x = \{z\} \wedge z \in Y) \}; Y \text{ ranges over subsets of } X: Y \in F \}$.

Let us consider X , let L be a continuous complete non empty poset, and let f be a function from the fixed ultrafilters of X into the carrier of L . The extension of f to homomorphism is a map from $(\text{Filt}(2_{\subseteq}^X), \subseteq)$ into L and is defined by the condition (Def. 3).

(Def. 3) Let F_1 be an element of $(\text{Filt}(2_{\subseteq}^X), \subseteq)$. Then (the extension of f to homomorphism)(F_1) = $\bigsqcup_L \{ \bigcap_L \{ f(\uparrow x) : \bigvee_z (x = \{z\} \wedge z \in Y) \}; Y \text{ ranges over subsets of } X: Y \in F_1 \}$.

We now state two propositions:

(12) Let L be a continuous complete non empty poset and f be a function from the fixed ultrafilters of X into the carrier of L . Then the extension of f to homomorphism is monotone.

(13) Let L be a continuous complete non empty poset and f be a function from the fixed ultrafilters of X into the carrier of L . Then (the extension of f to homomorphism)($\top_{(\text{Filt}(2_{\subseteq}^X), \subseteq)}$) = \top_L .

Let us consider X , let L be a continuous complete non empty poset, and let f be a function from the fixed ultrafilters of X into the carrier of L . One can verify that the extension of f to homomorphism is directed-sups-preserving.

Let us consider X , let L be a continuous complete non empty poset, and let f be a function from the fixed ultrafilters of X into the carrier of L . One can check that the extension of f to homomorphism is infs-preserving.

We now state several propositions:

(14) Let L be a continuous complete non empty poset and f be a function from the fixed ultrafilters of X into the carrier of L . Then (the extension of f to homomorphism)|(the fixed ultrafilters of X) = f .

(15) Let L be a continuous complete non empty poset, f be a function from the fixed ultrafilters of X into the carrier of L , and h be a CLHomomorphism of $(\text{Filt}(2_{\subseteq}^X), \subseteq), L$. Suppose $h \upharpoonright$ (the fixed ultrafilters of X) = f . Then h = the extension of f to homomorphism.

- (16) The fixed ultrafilters of X is a set of free generators of $\langle \text{Filt}(2_{\underline{C}}^X), \subseteq \rangle$.
- (17) Let L, M be continuous complete lattices and F, G be sets. Suppose F is a set of free generators of L and G is a set of free generators of M and $\overline{F} = \overline{G}$. Then L and M are isomorphic.
- (18) Let L be a continuous complete lattice and G be a set. Suppose G is a set of free generators of L and $\overline{G} = \overline{X}$. Then L and $\langle \text{Filt}(2_{\underline{C}}^X), \subseteq \rangle$ are isomorphic.

REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/card_1.html.
- [2] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.
- [3] Grzegorz Bancerek. Cartesian product of functions. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/funct_6.html.
- [4] Grzegorz Bancerek. Complete lattices. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/lattice3.html>.
- [5] Grzegorz Bancerek. Bounds in posets and relational substructures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_0.html.
- [6] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_0.html.
- [7] Grzegorz Bancerek. The "way-below" relation. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_3.html.
- [8] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [9] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [10] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zmisc_1.html.
- [11] Czesław Byliński. Galois connections. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_1.html.
- [12] Mariusz Giero. More on products of many sorted algebras. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/pralg_3.html.
- [13] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. *A Compendium of Continuous Lattices*. Springer-Verlag, Berlin, Heidelberg, New York, 1980.
- [14] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_1.html.
- [15] Robert Milewski. Completely-irreducible elements. *Journal of Formalized Mathematics*, 10, 1998. <http://mizar.org/JFM/Vol10/waybell6.html>.
- [16] Michał Muzalewski. Categories of groups. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/grcat_1.html.
- [17] Yatsuka Nakamura and Andrzej Trybulec. A mathematical model of CPU. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/ami_1.html.
- [18] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/setfam_1.html.
- [19] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.
- [20] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [21] Andrzej Trybulec. Function domains and Fränkel operator. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/fraenkel.html>.
- [22] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/pboole.html>.
- [23] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_1.html.
- [24] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/orders_1.html.
- [25] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [26] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

- [27] Mariusz Żynel and Czesław Byliński. Properties of relational structures, posets, lattices and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_2.html.

Received July 21, 1998

Published January 2, 2004
