

# Double Sequences and Iterated Limits in Regular Space

Roland Coghetto  
Rue de la Brasserie 5  
7100 La Louvière, Belgium

**Summary.** First, we define in Mizar [5], the Cartesian product of two filters bases and the Cartesian product of two filters. After comparing the product of two Fréchet filters on  $\mathbb{N}$  ( $\mathcal{F}_1$ ) with the Fréchet filter on  $\mathbb{N} \times \mathbb{N}$  ( $\mathcal{F}_2$ ), we compare  $\lim_{\mathcal{F}_1}$  and  $\lim_{\mathcal{F}_2}$  for all double sequences in a non empty topological space.

Endou, Okazaki and Shidama formalized in [14] the “convergence in Pringsheim’s sense” for double sequence of real numbers. We show some basic correspondences between the  $p$ -convergence and the filter convergence in a topological space. Then we formalize that the double sequence  $(x_{m,n} = \frac{1}{m+1})_{(m,n)} \in \mathbb{N} \times \mathbb{N}$  converges in “Pringsheim’s sense” but not in Fréchet filter on  $\mathbb{N} \times \mathbb{N}$  sense.

In the next section, we generalize some definitions: “is convergent in the first coordinate”, “is convergent in the second coordinate”, “the *lim* in the first coordinate of”, “the *lim* in the second coordinate of” according to [14], in Hausdorff space.

Finally, we generalize two theorems: (3) and (4) from [14] in the case of double sequences and we formalize the “iterated limit” theorem (“Double limit” [7], p. 81, par. 8.5 “*Double limite*” [6] (TG I,57)), all in regular space. We were inspired by the exercises (2.11.4), (2.17.5) [17] and the corrections B.10 [18].

MSC: 54A20 40A05 40B05 03B35

Keywords: filter; double limits; Pringsheim convergence; iterated limits; regular space

MML identifier: CARDFIL4, version: 8.1.05 5.37.1275

## 1. PRELIMINARIES

From now on  $x$  denotes an object,  $X, Y, Z$  denote sets,  $i, j, k, l, m, n$  denote natural numbers,  $r, s$  denote real numbers,  $n_1$  denotes an element of the ordered  $\mathbb{N}$ , and  $A$  denotes a subset of  $\mathbb{N} \times \mathbb{N}$ .

Now we state the propositions:

- (1) Let us consider a finite subset  $W$  of  $X$ . If  $X \setminus W \subseteq Z$ , then  $X \setminus Z$  is finite.
- (2) If  $Z \subseteq X$  and  $X \setminus Z$  is finite, then there exists a finite subset  $W$  of  $X$  such that  $X \setminus W = Z$ .
- (3) Let us consider sets  $X_1, X_2$ , a family  $S_1$  of subsets of  $X_1$ , and a family  $S_2$  of subsets of  $X_2$ . Then  $\{s, \text{ where } s \text{ is a subset of } X_1 \times X_2 : \text{ there exist sets } s_1, s_2 \text{ such that } s_1 \in S_1 \text{ and } s_2 \in S_2 \text{ and } s = s_1 \times s_2\}$  is a family of subsets of  $X_1 \times X_2$ .
- (4) If  $x \in X \times Y$ , then  $x$  is pair.
- (5) If  $0 < r$ , then there exists  $m$  such that  $m$  is not zero and  $\frac{1}{m} < r$ .
- (6) Let us consider points  $x, y$  of the metric space of real numbers. Then there exist real numbers  $x_1, y_1$  such that
  - (i)  $x = x_1$ , and
  - (ii)  $y = y_1$ , and
  - (iii)  $\rho(x, y) = \rho_{\mathbb{R}}(x, y)$ , and
  - (iv)  $\rho(x, y) = \rho^1(\langle x \rangle, \langle y \rangle)$ , and
  - (v)  $\rho(x, y) = |x_1 - y_1|$ .
- (7) Let us consider points  $x, y$  of  $(\mathcal{E}^1)_{\text{top}}$ . Then there exist points  $x_2, y_2$  of the metric space of real numbers and there exist real numbers  $x_1, y_1$  such that  $x_2 = x_1$  and  $y_2 = y_1$  and  $x = \langle x_1 \rangle$  and  $y = \langle y_1 \rangle$  and  $\rho(x_2, y_2) = \rho_{\mathbb{R}}(x_1, y_1)$  and  $\rho(x_2, y_2) = \rho^1(\langle x_1 \rangle, \langle y_1 \rangle)$  and  $\rho(x_2, y_2) = |x_1 - y_1|$ .
- (8) Let us consider points  $x, y$  of  $\mathcal{E}^1$ , and real numbers  $r, s$ . If  $x = \langle r \rangle$  and  $y = \langle s \rangle$ , then  $\rho(x, y) = |r - s|$ . The theorem is a consequence of (7).

One can check that  $\mathbb{N} \times \mathbb{N}$  is countable and  $\mathbb{N} \times \mathbb{N}$  is denumerable.

Now we state the propositions:

- (9) the set of all  $\langle 0, n \rangle$  where  $n$  is a natural number is infinite.  
 PROOF: Define  $\mathcal{F}(\text{object}) = \langle 0, \$1 \rangle$ . Consider  $f$  being a function such that  $\text{dom } f = \mathbb{N}$  and for every object  $x$  such that  $x \in \mathbb{N}$  holds  $f(x) = \mathcal{F}(x)$  from [9, Sch. 3].  $f$  is one-to-one.  $\text{rng } f =$  the set of all  $\langle 0, n \rangle$  where  $n$  is a natural number by [9, (3)].  $\square$
- (10) If  $i \leq k$  and  $j \leq l$ , then  $\mathbb{Z}_i \times \mathbb{Z}_j \subseteq \mathbb{Z}_k \times \mathbb{Z}_l$ .
- (11)  $(\mathbb{N} \setminus \mathbb{Z}_m) \times (\mathbb{N} \setminus \mathbb{Z}_n) \subseteq \mathbb{N} \times \mathbb{N} \setminus \mathbb{Z}_m \times \mathbb{Z}_n$ .
- (12) If  $n = n_1$  and  $n \leq m$ , then  $m \in \uparrow n_1$ .
- (13) If  $n = n_1$  and  $m \in \uparrow n_1$ , then  $n \leq m$ .
- (14) If  $n = n_1$ , then  $\uparrow n_1 = \mathbb{N} \setminus \mathbb{Z}_n$ .

PROOF:  $\uparrow n_1 \subseteq \mathbb{N} \setminus \mathbb{Z}_n$  by [12, (50)], (13), [1, (44)].  $\mathbb{N} \setminus \mathbb{Z}_n \subseteq \uparrow n_1$  by [1, (44)], [12, (50)].  $\square$

- (15)  $\pi_1(A) = \{x, \text{ where } x \text{ is an element of } \mathbb{N} : \text{ there exists an element } y \text{ of } \mathbb{N} \text{ such that } \langle x, y \rangle \in A\}$ .
- (16)  $\pi_2(A) = \{y, \text{ where } y \text{ is an element of } \mathbb{N} : \text{ there exists an element } x \text{ of } \mathbb{N} \text{ such that } \langle x, y \rangle \in A\}$ .
- (17) Let us consider a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$ . Then there exists  $m$  and there exists  $n$  such that  $A \subseteq \mathbb{Z}_m \times \mathbb{Z}_n$ . The theorem is a consequence of (15) and (16).
- (18) Let us consider a non empty set  $X$ . Then every filter of  $X$  is a proper filter of  $2_{\subseteq}^X$ .
- (19) Let us consider a non empty set  $X$ , and a filter  $\mathcal{F}$  of  $X$ . Then there exists a filter base  $\mathcal{B}$  of  $X$  such that
  - (i)  $\mathcal{B} = \mathcal{F}$ , and
  - (ii)  $[\mathcal{B}] = \mathcal{F}$ .
- (20) Let us consider a non empty topological space  $T$ , and a filter  $\mathcal{F}$  of the carrier of  $T$ . If  $x \in \text{LimFilter}(\mathcal{F})$ , then  $x$  is a cluster point of  $\mathcal{F}, T$ .
- (21) Let us consider an element  $B$  of the base of Frechet filter. Then there exists  $n$  such that  $B = \mathbb{N} \setminus \mathbb{Z}_n$ . The theorem is a consequence of (14).
- (22) Let us consider a subset  $B$  of  $\mathbb{N}$ . Suppose  $B = \mathbb{N} \setminus \mathbb{Z}_n$ . Then  $B$  is an element of the base of Frechet filter. The theorem is a consequence of (14).

## 2. CARTESIAN PRODUCT OF TWO FILTERS

From now on  $X, Y, X_1, X_2$  denote non empty sets,  $\mathcal{A}_1, \mathcal{B}_1$  denote filter bases of  $X_1$ ,  $\mathcal{A}_2, \mathcal{B}_2$  denote filter bases of  $X_2$ ,  $\mathcal{F}_1$  denotes a filter of  $X_1$ ,  $\mathcal{F}_2$  denotes a filter of  $X_2$ ,  $\mathcal{B}_3$  denotes a generalized basis of  $\mathcal{F}_1$ .

Let  $X_1, X_2$  be non empty sets,  $\mathcal{B}_1$  be a filter base of  $X_1$ , and  $\mathcal{B}_2$  be a filter base of  $X_2$ . The functor  $\mathcal{B}_1 \times \mathcal{B}_2$  yielding a filter base of  $X_1 \times X_2$  is defined by the term

(Def. 1) the set of all  $B_1 \times B_2$  where  $B_1$  is an element of  $\mathcal{B}_1$ ,  $B_2$  is an element of  $\mathcal{B}_2$ .

Now we state the propositions:

- (23) Suppose  $\mathcal{F}_1 = [\mathcal{B}_1)$  and  $\mathcal{F}_1 = [\mathcal{A}_1)$  and  $\mathcal{F}_2 = [\mathcal{B}_2)$  and  $\mathcal{F}_2 = [\mathcal{A}_2)$ . Then  $[\mathcal{B}_1 \times \mathcal{B}_2) = [\mathcal{A}_1 \times \mathcal{A}_2)$ .
- (24) If  $\mathcal{B}_3 = \mathcal{B}_1$ , then  $[\mathcal{B}_1] = \mathcal{F}_1$ .

(25) There exists  $\mathcal{B}_1$  such that  $[\mathcal{B}_1] = \mathcal{F}_1$ . The theorem is a consequence of (24).

Let  $X_1, X_2$  be non empty sets,  $\mathcal{F}_1$  be a filter of  $X_1$ , and  $\mathcal{F}_2$  be a filter of  $X_2$ . The functor  $\langle \mathcal{F}_1, \mathcal{F}_2 \rangle$  yielding a filter of  $X_1 \times X_2$  is defined by

(Def. 2) there exists a filter base  $\mathcal{B}_1$  of  $X_1$  and there exists a filter base  $\mathcal{B}_2$  of  $X_2$  such that  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$  and  $it = [\mathcal{B}_1 \times \mathcal{B}_2]$ .

Let  $\mathcal{B}_1$  be a generalized basis of  $\mathcal{F}_1$  and  $\mathcal{B}_2$  be a generalized basis of  $\mathcal{F}_2$ . The functor  $\mathcal{B}_1 \times \mathcal{B}_2$  yielding a generalized basis of  $\langle \mathcal{F}_1, \mathcal{F}_2 \rangle$  is defined by

(Def. 3) there exists a filter base  $\mathcal{B}_3$  of  $X_1$  and there exists a filter base  $\mathcal{B}_4$  of  $X_2$  such that  $\mathcal{B}_1 = \mathcal{B}_3$  and  $\mathcal{B}_2 = \mathcal{B}_4$  and  $it = \mathcal{B}_3 \times \mathcal{B}_4$ .

Let  $n$  be a natural number. The functor  $\uparrow^2(n)$  yielding a subset of  $\mathbb{N} \times \mathbb{N}$  is defined by

(Def. 4) for every element  $x$  of  $\mathbb{N} \times \mathbb{N}$ ,  $x \in it$  iff there exist natural numbers  $n_1, n_2$  such that  $n_1 = (x)_1$  and  $n_2 = (x)_2$  and  $n \leq n_1$  and  $n \leq n_2$ .

Now we state the proposition:

(26)  $\langle n, n \rangle \in \uparrow^2(n)$ .

Let us consider  $n$ . One can check that  $\uparrow^2(n)$  is non empty.

Now we state the propositions:

(27) If  $\langle i, j \rangle \in \uparrow^2(n)$ , then  $\langle i + k, j \rangle, \langle i, j + l \rangle \in \uparrow^2(n)$ .

(28)  $\uparrow^2(n)$  is an infinite subset of  $\mathbb{N} \times \mathbb{N}$ . The theorem is a consequence of (17).

(29) If  $n_1 = n$ , then  $\uparrow^2(n) = \uparrow n_1 \times \uparrow n_1$ . The theorem is a consequence of (12) and (13).

(30) If  $m = n - 1$ , then  $\uparrow^2(n) \subseteq \mathbb{N} \times \mathbb{N} \setminus \text{Seg } m \times \text{Seg } m$ .

PROOF: Reconsider  $y = x$  as an element of  $\mathbb{N} \times \mathbb{N}$ . Consider  $n_1, n_2$  being natural numbers such that  $n_1 = (y)_1$  and  $n_2 = (y)_2$  and  $n \leq n_1$  and  $n \leq n_2$ .  $x \notin \text{Seg } m \times \text{Seg } m$  by [3, (1)].  $\square$

(31)  $\uparrow^2(n) \subseteq \mathbb{N} \times \mathbb{N} \setminus \mathbb{Z}_n \times \mathbb{Z}_n$ .

PROOF: Reconsider  $y = x$  as an element of  $\mathbb{N} \times \mathbb{N}$ . Consider  $n_1, n_2$  being natural numbers such that  $n_1 = (y)_1$  and  $n_2 = (y)_2$  and  $n \leq n_1$  and  $n \leq n_2$ .  $x \notin \mathbb{Z}_n \times \mathbb{Z}_n$  by [16, (10)].  $\square$

(32)  $\uparrow^2(n) = (\mathbb{N} \setminus \mathbb{Z}_n) \times (\mathbb{N} \setminus \mathbb{Z}_n)$ . The theorem is a consequence of (14) and (29).

(33) There exists  $n$  such that  $\uparrow^2(n) \subseteq (\mathbb{N} \setminus \mathbb{Z}_i) \times (\mathbb{N} \setminus \mathbb{Z}_j)$ . The theorem is a consequence of (4).

(34) If  $n = \max(i, j)$ , then  $\uparrow^2(n) \subseteq (\uparrow^2(i)) \cap (\uparrow^2(j))$ .

Let  $n$  be a natural number. The functor  $\downarrow^2(n)$  yielding a subset of  $\mathbb{N} \times \mathbb{N}$  is defined by

(Def. 5) for every element  $x$  of  $\mathbb{N} \times \mathbb{N}$ ,  $x \in it$  iff there exist natural numbers  $n_1, n_2$  such that  $n_1 = (x)_1$  and  $n_2 = (x)_2$  and  $n_1 < n$  and  $n_2 < n$ .

Now we state the propositions:

(35)  $\downarrow^2(n) = \mathbb{Z}_n \times \mathbb{Z}_n$ .

PROOF:  $\downarrow^2(n) \subseteq \mathbb{Z}_n \times \mathbb{Z}_n$  by [1, (44)]. Consider  $y_2, y_1$  being objects such that  $y_2 \in \mathbb{Z}_n$  and  $y_1 \in \mathbb{Z}_n$  and  $x = \langle y_2, y_1 \rangle$ .  $\square$

(36) Let us consider a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$ . Then there exists  $n$  such that  $A \subseteq \downarrow^2(n)$ .

PROOF: Consider  $m, n$  such that  $A \subseteq \mathbb{Z}_m \times \mathbb{Z}_n$ . Reconsider  $m_1 = \max(m, n)$  as a natural number.  $A \subseteq \downarrow^2(m_1)$  by [1, (39)], [11, (96)], (35).  $\square$

(37)  $\downarrow^2(n)$  is a finite subset of  $\mathbb{N} \times \mathbb{N}$ . The theorem is a consequence of (35).

### 3. COMPARISON BETWEEN CARTESIAN PRODUCT OF FRECHET FILTER ON $\mathbb{N}$ AND THE FRECHET FILTER OF $\mathbb{N} \times \mathbb{N}$

Let us consider an element  $x$  of (the base of Frechet filter)  $\times$  (the base of Frechet filter). Now we state the propositions:

(38) There exists  $i$  and there exists  $j$  such that  $x = (\mathbb{N} \setminus \mathbb{Z}_i) \times (\mathbb{N} \setminus \mathbb{Z}_j)$ . The theorem is a consequence of (21).

(39) There exists  $n$  such that  $\uparrow^2(n) \subseteq x$ . The theorem is a consequence of (38) and (33).

(40) (The base of Frechet filter)  $\times$  (the base of Frechet filter) is a filter base of  $\mathbb{N} \times \mathbb{N}$ .

(41) There exists a generalized basis  $\mathcal{B}$  of  $\text{FrechetFilter}(\mathbb{N})$  such that

(i)  $\mathcal{B}$  = the base of Frechet filter, and

(ii)  $\mathcal{B} \times \mathcal{B}$  is a generalized basis of  $\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$ .

The functor  $\uparrow_{\mathbb{N}}^2$  yielding a filter base of  $\mathbb{N} \times \mathbb{N}$  is defined by the term

(Def. 6) the set of all  $\uparrow^2(n)$  where  $n$  is a natural number.

Now we state the propositions:

(42)  $\uparrow_{\mathbb{N}}^2$  and (the base of Frechet filter)  $\times$  (the base of Frechet filter) are equivalent generators. The theorem is a consequence of (22), (32), and (39).

(43)  $[(\text{the base of Frechet filter}) \times (\text{the base of Frechet filter})] = \langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$ . The theorem is a consequence of (41).

(44)  $[\uparrow_{\mathbb{N}}^2] = \langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$ .

(45)  $\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$  is finer than  $\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})$ .  
 The theorem is a consequence of (17), (11), (22), and (43).

(46) (i)  $\mathbb{N} \times \mathbb{N} \setminus$  the set of all  $\langle 0, n \rangle$  where  $n$  is a natural number  $\in \langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$ , and

(ii)  $\mathbb{N} \times \mathbb{N} \setminus$  the set of all  $\langle 0, n \rangle$  where  $n$  is a natural number  $\notin \text{FrechetFilter}(\mathbb{N} \times \mathbb{N})$ .

PROOF: Set  $X = \mathbb{N} \times \mathbb{N} \setminus$  the set of all  $\langle 0, n \rangle$  where  $n$  is a natural number.  $\uparrow^2(1) \subseteq X$  by (32), [1, (44)].  $X \notin \text{FrechetFilter}(\mathbb{N} \times \mathbb{N})$  by [12, (51)], [15, (5)], (9).  $\square$

(47)  $\text{FrechetFilter}(\mathbb{N} \times \mathbb{N}) \neq \langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$ .

#### 4. TOPOLOGICAL SPACE AND DOUBLE SEQUENCE

In the sequel  $T$  denotes a non empty topological space,  $s$  denotes a function from  $\mathbb{N} \times \mathbb{N}$  into the carrier of  $T$ ,  $M$  denotes a subset of the carrier of  $T$ , and  $\mathcal{F}_1, \mathcal{F}_2$  denote filters of the carrier of  $T$ . Now we state the propositions:

(48) If  $\mathcal{F}_2$  is finer than  $\mathcal{F}_1$ , then  $\text{LimFilter}(\mathcal{F}_1) \subseteq \text{LimFilter}(\mathcal{F}_2)$ .

(49) Let us consider a function  $f$  from  $X$  into  $Y$ , and filters  $\mathcal{F}_1, \mathcal{F}_2$  of  $X$ . Suppose  $\mathcal{F}_2$  is finer than  $\mathcal{F}_1$ . Then the image of filter  $\mathcal{F}_2$  under  $f$  is finer than the image of filter  $\mathcal{F}_1$  under  $f$ .

(50)  $s^{-1}(M) \in \text{FrechetFilter}(\mathbb{N} \times \mathbb{N})$  if and only if there exists a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$  such that  $s^{-1}(M) = \mathbb{N} \times \mathbb{N} \setminus A$ .

(51)  $s^{-1}(M) \in \langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$  if and only if there exists  $n$  such that  $\uparrow^2(n) \subseteq s^{-1}(M)$ . The theorem is a consequence of (43), (39), and (42).

(52) The image of filter  $\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})$  under  $s = \{M$ , where  $M$  is a subset of the carrier of  $T$  : there exists a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$  such that  $s^{-1}(M) = \mathbb{N} \times \mathbb{N} \setminus A\}$ . The theorem is a consequence of (50).

(53) The image of filter  $\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle$  under  $s = \{M$ , where  $M$  is a subset of the carrier of  $T$  : there exists a natural number  $n$  such that  $\uparrow^2(n) \subseteq s^{-1}(M)\}$ . The theorem is a consequence of (51).

Let us consider a point  $x$  of  $T$ . Now we state the propositions:

(54)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every neighbourhood  $A$  of  $x$ , there exists a finite subset  $B$  of  $\mathbb{N} \times \mathbb{N}$  such that  $s^{-1}(A) = \mathbb{N} \times \mathbb{N} \setminus B$ . The theorem is a consequence of (52).

(55)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every neighbourhood  $A$  of  $x$ ,  $\mathbb{N} \times \mathbb{N} \setminus s^{-1}(A)$  is finite. The theorem is a consequence of (54), (1), and (2).

- (56)  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every neighbourhood  $A$  of  $x$ , there exists a natural number  $n$  such that  $\uparrow^2(n) \subseteq s^{-1}(A)$ . The theorem is a consequence of (53).

Let us consider a point  $x$  of  $T$  and a generalized basis  $\mathcal{B}$  of BooleanFilter ToFilter(the neighborhood system of  $x$ ). Now we state the propositions:

- (57)  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists a natural number  $n$  such that  $\uparrow^2(n) \subseteq s^{-1}(B)$ . The theorem is a consequence of (56).
- (58)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$  such that  $s^{-1}(B) = \mathbb{N} \times \mathbb{N} \setminus A$ . The theorem is a consequence of (54), (1), and (55).
- (59)  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists a natural number  $n$  such that  $s^\circ(\uparrow^2(n)) \subseteq B$ . The theorem is a consequence of (57).

- (60)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists a finite subset  $A$  of  $\mathbb{N} \times \mathbb{N}$  such that  $s^\circ(\mathbb{N} \times \mathbb{N} \setminus A) \subseteq B$ .

PROOF: For every neighbourhood  $A$  of  $x$ ,  $\mathbb{N} \times \mathbb{N} \setminus s^{-1}(A)$  is finite by [4, (2)], [19, (143)], [9, (76)].  $\square$

- (61)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists  $n$  and there exists  $m$  such that  $s^\circ(\mathbb{N} \times \mathbb{N} \setminus \mathbb{Z}_n \times \mathbb{Z}_m) \subseteq B$ . The theorem is a consequence of (60) and (17).

- (62)  $x \in s^\circ(\uparrow^2(n))$  if and only if there exists  $i$  and there exists  $j$  such that  $n \leq i$  and  $n \leq j$  and  $x = s(i, j)$ .

- (63)  $x \in s^\circ(\mathbb{N} \times \mathbb{N} \setminus \mathbb{Z}_i \times \mathbb{Z}_j)$  if and only if there exist natural numbers  $n, m$  such that  $(i \leq n$  or  $j \leq m)$  and  $x = s(n, m)$ .

PROOF: Consider  $n, m$  being natural numbers such that  $i \leq n$  or  $j \leq m$  and  $x = s(n, m)$ .  $\langle n, m \rangle \notin \mathbb{Z}_i \times \mathbb{Z}_j$  by [1, (44)].  $\square$

Let us consider a point  $x$  of  $T$  and a generalized basis  $\mathcal{B}$  of BooleanFilter ToFilter(the neighborhood system of  $x$ ). Now we state the propositions:

- (64)  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $s(n_1, n_2) \in B$ . The theorem is a consequence of (62) and (59).

- (65)  $x \in \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s$  if and only if for every element  $B$  of  $\mathcal{B}$ , there exists  $i$  and there exists  $j$  such that for every  $m$  and  $n$  such that  $i \leq m$  or  $j \leq n$  holds  $s(m, n) \in B$ . The theorem is a consequence of (61).

- (66)  $\lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} s \subseteq \lim_{[\uparrow^2_{\mathbb{N}}]} s$ . The theorem is a consequence of (42), (43), (45), (48), and (49).

## 5. METRIC SPACE AND DOUBLE SEQUENCE

Now we state the propositions:

- (67) Let us consider a non empty metric space  $M$ , a point  $p$  of  $M$ , a point  $x$  of  $M_{\text{top}}$ , and a function  $s$  from  $\mathbb{N} \times \mathbb{N}$  into  $M_{\text{top}}$ . Suppose  $x = p$ . Then  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $s(n_1, n_2) \in \{q, \text{ where } q \text{ is a point of } M : \rho(p, q) < \frac{1}{m}\}$ .

PROOF:  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  iff for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $s(n_1, n_2) \in \{q, \text{ where } q \text{ is a point of } M : \rho(p, q) < \frac{1}{m}\}$  by [13, (6)], (64).  $\square$

- (68) Let us consider a non empty metric space  $M$ , a point  $p$  of  $M$ , a point  $x$  of  $M_{\text{top}}$ , a function  $s$  from  $\mathbb{N} \times \mathbb{N}$  into  $M_{\text{top}}$ , and a function  $s_2$  from  $\mathbb{N} \times \mathbb{N}$  into  $M$ . Suppose  $x = p$  and  $s = s_2$ . Then  $x \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} s$  if and only if for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $s_2(n_1, n_2) \in \{q, \text{ where } q \text{ is a point of } M : \rho(p, q) < \frac{1}{m}\}$ .

## 6. ONE-DIMENSIONAL EUCLIDEAN METRIC SPACE AND DOUBLE SEQUENCE

In the sequel  $R$  denotes a function from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}$ .

Now we state the proposition:

- (69) Let us consider a point  $x$  of  $(\mathcal{E}^1)_{\text{top}}$ , a point  $y$  of  $\mathcal{E}^1$ , a generalized basis  $\mathcal{B}$  of  $\text{BooleanFilterToFilter}(\text{the neighborhood system of } x)$ , and an element  $b$  of  $\mathcal{B}$ . Suppose  $x = y$  and  $\mathcal{B} = \text{Balls } x$ . Then there exists a natural number  $n$  such that  $b = \{q, \text{ where } q \text{ is an element of } \mathcal{E}^1 : \rho(y, q) < \frac{1}{n}\}$ .

Let  $s$  be a function from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}$ . The functor  $\# s$  yielding a function from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}^1$  is defined by the term

(Def. 7)  $s$ .

Now we state the propositions:

- (70) Let us consider a function  $s$  from  $\mathbb{N} \times \mathbb{N}$  into  $(\mathcal{E}^1)_{\text{top}}$ , and a point  $y$  of  $\mathcal{E}^1$ . Then  $s^\circ(\uparrow^2(n)) \subseteq \{q, \text{ where } q \text{ is an element of } \mathcal{E}^1 : \rho(y, q) < \frac{1}{m}\}$  if and only if for every object  $x$  such that  $x \in s^\circ(\uparrow^2(n))$  there exist real numbers  $r_1, r_2$  such that  $x = \langle r_1 \rangle$  and  $y = \langle r_2 \rangle$  and  $|r_2 - r_1| < \frac{1}{m}$ . The theorem is a consequence of (8).

- (71)  $r \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R$  if and only if for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every

natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $|R(n_1, n_2) - r| < \frac{1}{m}$ .

PROOF: Reconsider  $p = r$  as a point of the metric space of real numbers. for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $R(n_1, n_2) \in \{q, \text{ where } q \text{ is a point of the metric space of real numbers} : \rho(p, q) < \frac{1}{m}\}$  iff for every non zero natural number  $m$ , there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $|R(n_1, n_2) - r| < \frac{1}{m}$  by (6), [8, (60)].  $\square$

### 7. BASIC RELATIONS CONVERGENCE IN PRINGSHEIM'S SENSE AND FILTER CONVERGENCE

Now we state the propositions:

- (72) Suppose  $\lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R \neq \emptyset$ . Then there exists a real number  $x$  such that  $\lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R = \{x\}$ .
- (73) If  $R$  is P-convergent, then  $\text{P-lim } R \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R$ . The theorem is a consequence of (71).
- (74)  $R$  is P-convergent if and only if  $\lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R \neq \emptyset$ . The theorem is a consequence of (71) and (5).
- (75) Suppose  $R$  is P-convergent. Then  $\{\text{P-lim } R\} = \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R$ . The theorem is a consequence of (73) and (72).
- (76) Suppose  $\lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R$  is not empty. Then
  - (i)  $R$  is P-convergent, and
  - (ii)  $\{\text{P-lim } R\} = \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# R$ .

### 8. EXAMPLE: DOUBLE SEQUENCE CONVERGES IN PRINGSHEIM'S SENSE BUT NOT IN FRECHET FILTER OF $\mathbb{N} \times \mathbb{N}$ SENSE

The functor  $\text{DbSeq-ex1}$  yielding a function from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}$  is defined by (Def. 8) for every natural numbers  $m, n$ ,  $it(m, n) = \frac{1}{m+1}$ .

Now we state the propositions:

- (77) Let us consider a non zero natural number  $m$ . Then there exists a natural number  $n$  such that for every natural numbers  $n_1, n_2$  such that  $n \leq n_1$  and  $n \leq n_2$  holds  $|(\text{DbSeq-ex1})(n_1, n_2) - 0| < \frac{1}{m}$ .
- (78)  $0 \in \lim_{\langle \text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}) \rangle} \# \text{DbSeq-ex1}$ .

- (79)  $\lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} \# \text{DblSeq-ex1} = \emptyset$ . The theorem is a consequence of (66), (42), (43), (72), (78), and (65).
- (80)  $\lim_{(\text{FrechetFilter}(\mathbb{N}), \text{FrechetFilter}(\mathbb{N}))} \# \text{DblSeq-ex1} \neq \lim_{\text{FrechetFilter}(\mathbb{N} \times \mathbb{N})} \# \text{DblSeq-ex1}$ .

9. CORRESPONDENCE WITH SOME DEFINITIONS FROM [14]

Let  $X_1, X_2$  be non empty sets,  $\mathcal{F}_1$  be a filter of  $X_1$ ,  $Y$  be a Hausdorff, non empty topological space, and  $f$  be a function from  $X_1 \times X_2$  into  $Y$ . Assume for every element  $x$  of  $X_2$ ,  $\lim_{\mathcal{F}_1} \text{curry}'(f, x) \neq \emptyset$ . The functor  $\lim_1(f, \mathcal{F}_1)$  yielding a function from  $X_2$  into  $Y$  is defined by

(Def. 9) for every element  $x$  of  $X_2$ ,  $\{it(x)\} = \lim_{\mathcal{F}_1} \text{curry}'(f, x)$ .

Let  $\mathcal{F}_2$  be a filter of  $X_2$ . Assume for every element  $x$  of  $X_1$ ,  $\lim_{\mathcal{F}_2} \text{curry}(f, x) \neq \emptyset$ . The functor  $\lim_2(f, \mathcal{F}_2)$  yielding a function from  $X_1$  into  $Y$  is defined by

(Def. 10) for every element  $x$  of  $X_1$ ,  $\{it(x)\} = \lim_{\mathcal{F}_2} \text{curry}(f, x)$ .

Now we state the propositions:

- (81) Every function from  $X$  into  $\mathbb{R}$  is a function from  $X$  into  $\mathbb{R}^1$ .
- (82) Every sequence of  $\mathbb{R}$  is a function from  $\mathbb{N}$  into  $\mathbb{R}^1$ .

From now on  $f$  denotes a function from  $\Omega_{\text{the ordered } \mathbb{N}}$  into  $\mathbb{R}^1$  and  $s_1$  denotes a function from  $\mathbb{N}$  into  $\mathbb{R}$ .

Now we state the propositions:

- (83) Suppose  $f = s_1$  and  $\text{LimF}(f) \neq \emptyset$ . Then
  - (i)  $s_1$  is convergent, and
  - (ii) there exists a real number  $z$  such that  $z \in \text{LimF}(f)$  and for every real number  $p$  such that  $0 < p$  there exists a natural number  $n$  such that for every natural number  $m$  such that  $n \leq m$  holds  $|s_1(m) - z| < p$ .

PROOF: Consider  $x$  being an object such that  $x \in \text{LimF}(f)$ . Reconsider  $y = x$  as a point of (the metric space of real numbers)<sub>top</sub>. Reconsider  $z = y$  as a real number. Consider  $y_1$  being a point of the metric space of real numbers such that  $y_1 = y$  and  $\text{Balls } y = \{\text{Ball}(y_1, \frac{1}{n})\}$ , where  $n$  is a natural number :  $n \neq 0$ . For every real number  $p$  such that  $0 < p$  there exists a natural number  $n$  such that for every natural number  $m$  such that  $n \leq m$  holds  $|s_1(m) - z| < p$  by (5), [12, (84), (50)], [2, (18)].  $\square$

- (84) If  $f = s_1$  and  $\text{LimF}(f) \neq \emptyset$ , then  $\text{LimF}(f) = \{\lim s_1\}$ .

PROOF: Consider  $x$  being an object such that  $x \in \text{LimF}(f)$ . Consider  $u$  being an object such that  $\text{LimF}(f) = \{u\}$ .  $\text{LimF}(f) = \{\lim s_1\}$  by (83), [11, (3)].  $\square$

- (85) Let us consider a function  $f$  from  $\Omega_\alpha$  into  $T$ , and a sequence  $s$  of  $T$ . If  $f = s$ , then  $\text{LimF}(f) = \text{LimF}(s)$ , where  $\alpha$  is the ordered  $\mathbb{N}$ .
- (86) Let us consider a function  $f$  from  $\Omega_\alpha$  into  $T$ , and a function  $g$  from  $\mathbb{N}$  into  $T$ . If  $f = g$ , then  $\text{LimF}(f) = \text{LimF}(g)$ , where  $\alpha$  is the ordered  $\mathbb{N}$ .
- (87) Let us consider a function  $f$  from  $\mathbb{N}$  into  $\mathbb{R}^1$ . Suppose  $f = s_1$  and  $\text{LimF}(f) \neq \emptyset$ . Then  $\text{LimF}(f) = \{\lim s_1\}$ . The theorem is a consequence of (84).
- (88) for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}'(\# R, x) \neq \emptyset$  if and only if  $R$  is convergent in the first coordinate. The theorem is a consequence of (5).
- (89) for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}(\# R, x) \neq \emptyset$  if and only if  $R$  is convergent in the second coordinate. The theorem is a consequence of (5).

Let us consider an element  $t$  of  $\mathbb{N}$ , a function  $f$  from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}^1$ , and a function  $s_1$  from  $\mathbb{N} \times \mathbb{N}$  into  $\mathbb{R}$ . Now we state the propositions:

- (90) Suppose  $f = s_1$  and for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}(f, x) \neq \emptyset$ . Then  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}(f, t) = \{\lim \text{curry}(s_1, t)\}$ . The theorem is a consequence of (87).
- (91) Suppose  $f = s_1$  and for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}'(f, x) \neq \emptyset$ . Then  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}'(f, t) = \{\lim \text{curry}'(s_1, t)\}$ . The theorem is a consequence of (87).
- (92) Let us consider a Hausdorff, non empty topological space  $Y$ , and a function  $f$  from  $\mathbb{N} \times \mathbb{N}$  into  $Y$ . Suppose for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}'(f, x) \neq \emptyset$  and  $f = R$  and  $Y = \mathbb{R}^1$ . Then  $\lim_1(f, \text{FrechetFilter}(\mathbb{N})) =$  the lim in the first coordinate of  $R$ . The theorem is a consequence of (91).
- (93) Let us consider a non empty, Hausdorff topological space  $Y$ , and a function  $f$  from  $\mathbb{N} \times \mathbb{N}$  into  $Y$ . Suppose for every element  $x$  of  $\mathbb{N}$ ,  $\lim_{\text{FrechetFilter}(\mathbb{N})} \text{curry}(f, x) \neq \emptyset$  and  $f = R$  and  $Y = \mathbb{R}^1$ . Then  $\lim_2(f, \text{FrechetFilter}(\mathbb{N})) =$  the lim in the second coordinate of  $R$ . The theorem is a consequence of (90).

## 10. REGULAR SPACE, DOUBLE LIMIT AND ITERATED LIMIT

From now on  $Y$  denotes a non empty topological space,  $x$  denotes a point of  $Y$ , and  $f$  denotes a function from  $X_1 \times X_2$  into  $Y$ .

Now we state the proposition:

- (94) Suppose  $x \in \lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f$  and  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$ . Let us consider a subset  $V$  of  $Y$ . Suppose  $V$  is open and  $x \in V$ . Then there exists an ele-

ment  $B_1$  of  $\mathcal{B}_1$  and there exists an element  $B_2$  of  $\mathcal{B}_2$  such that  $f^\circ(B_1 \times B_2) \subseteq V$ .

Let us consider a neighbourhood  $U$  of  $x$ . Now we state the propositions:

- (95) Suppose  $x \in \lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f$  and  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$ . Then suppose  $U$  is closed. Then there exists an element  $B_1$  of  $\mathcal{B}_1$  and there exists an element  $B_2$  of  $\mathcal{B}_2$  such that  $f^\circ(B_1 \times B_2) \subseteq \text{Int } U$ .
- (96) Suppose  $x \in \lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f$  and  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$ . Then suppose  $U$  is closed. Then there exists an element  $B_1$  of  $\mathcal{B}_1$  and there exists an element  $B_2$  of  $\mathcal{B}_2$  such that for every element  $y$  of  $B_1$ ,  $f^\circ(\{y\} \times B_2) \subseteq \text{Int } U$ . The theorem is a consequence of (95).
- (97) Suppose  $x \in \lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f$  and  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$ . Then suppose  $U$  is closed. Then there exists an element  $B_1$  of  $\mathcal{B}_1$  and there exists an element  $B_2$  of  $\mathcal{B}_2$  such that for every element  $z$  of  $X_1$  for every element  $y$  of  $Y$  such that  $z \in B_1$  and  $y \in \lim_{\mathcal{F}_2} \text{curry}(f, z)$  holds  $y \in \overline{\text{Int } U}$ .

PROOF: Consider  $B_1$  being an element of  $\mathcal{B}_1$ ,  $B_2$  being an element of  $\mathcal{B}_2$  such that  $f^\circ(B_1 \times B_2) \subseteq \text{Int } U$ . For every element  $y$  of  $B_1$ ,  $f^\circ(\{y\} \times B_2) \subseteq \text{Int } U$  by [11, (95)], [19, (125)]. For every element  $z$  of  $B_1$  and for every element  $y$  of  $Y$  such that  $y \in \lim_{\mathcal{F}_2} \text{curry}(f, z)$  holds the image of filter  $\mathcal{F}_2$  under  $\text{curry}(f, z)$  is a proper filter of  $2_{\subseteq}^{\Omega_Y}$  and  $\text{Int } U \in$  the image of filter  $\mathcal{F}_2$  under  $\text{curry}(f, z)$  and  $y$  is a cluster point of the image of filter  $\mathcal{F}_2$  under  $\text{curry}(f, z)$ ,  $Y$  by (18), [19, (132)], [10, (95)], (20). For every element  $z$  of  $B_1$  and for every element  $y$  of  $Y$  such that  $y \in \lim_{\mathcal{F}_2} \text{curry}(f, z)$  holds  $y \in \overline{\text{Int } U}$  by [4, (25)].  $\square$

- (98) Suppose  $x \in \lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f$  and  $[\mathcal{B}_1] = \mathcal{F}_1$  and  $[\mathcal{B}_2] = \mathcal{F}_2$ . Then suppose  $U$  is closed. Then there exists an element  $B_1$  of  $\mathcal{B}_1$  and there exists an element  $B_2$  of  $\mathcal{B}_2$  such that for every element  $z$  of  $X_2$  for every element  $y$  of  $Y$  such that  $z \in B_2$  and  $y \in \lim_{\mathcal{F}_1} \text{curry}'(f, z)$  holds  $y \in \overline{\text{Int } U}$ .

PROOF: Consider  $B_1$  being an element of  $\mathcal{B}_1$ ,  $B_2$  being an element of  $\mathcal{B}_2$  such that  $f^\circ(B_1 \times B_2) \subseteq \text{Int } U$ . For every element  $y$  of  $B_2$ ,  $f^\circ(B_1 \times \{y\}) \subseteq \text{Int } U$  by [11, (95)], [19, (125)]. For every element  $z$  of  $B_2$  and for every element  $y$  of  $Y$  such that  $y \in \lim_{\mathcal{F}_1} \text{curry}'(f, z)$  holds the image of filter  $\mathcal{F}_1$  under  $\text{curry}'(f, z)$  is a proper filter of  $2_{\subseteq}^{\Omega_Y}$  and  $\text{Int } U \in$  the image of filter  $\mathcal{F}_1$  under  $\text{curry}'(f, z)$  and  $y$  is a cluster point of the image of filter  $\mathcal{F}_1$  under  $\text{curry}'(f, z)$ ,  $Y$  by (18), [19, (132)], [10, (95)], (20). For every element  $z$  of  $B_2$  and for every element  $y$  of  $Y$  such that  $y \in \lim_{\mathcal{F}_1} \text{curry}'(f, z)$  holds  $y \in \overline{\text{Int } U}$  by [4, (25)].  $\square$

Let us consider a Hausdorff, regular, non empty topological space  $Y$  and a function  $f$  from  $X_1 \times X_2$  into  $Y$ . Now we state the propositions:

- (99) Suppose for every element  $x$  of  $X_2$ ,  $\lim_{\mathcal{F}_1} \text{curry}'(f, x) \neq \emptyset$ . Then  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle}$

$f \subseteq \lim_{\mathcal{F}_2} \lim_1(f, \mathcal{F}_1)$ . The theorem is a consequence of (19) and (98).

- (100) Suppose for every element  $x$  of  $X_1$ ,  $\lim_{\mathcal{F}_2} \text{curry}(f, x) \neq \emptyset$ . Then  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f \subseteq \lim_{\mathcal{F}_1} \lim_2(f, \mathcal{F}_2)$ . The theorem is a consequence of (19) and (97).

Let us consider non empty sets  $X_1$ ,  $X_2$ , a filter  $\mathcal{F}_1$  of  $X_1$ , a filter  $\mathcal{F}_2$  of  $X_2$ , a Hausdorff, regular, non empty topological space  $Y$ , and a function  $f$  from  $X_1 \times X_2$  into  $Y$ . Now we state the propositions:

- (101) Suppose  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f \neq \emptyset$  and for every element  $x$  of  $X_1$ ,  $\lim_{\mathcal{F}_2} \text{curry}(f, x) \neq \emptyset$ . Then  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f = \lim_{\mathcal{F}_1} \lim_2(f, \mathcal{F}_2)$ . The theorem is a consequence of (100).
- (102) Suppose  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f \neq \emptyset$  and for every element  $x$  of  $X_2$ ,  $\lim_{\mathcal{F}_1} \text{curry}'(f, x) \neq \emptyset$ . Then  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f = \lim_{\mathcal{F}_2} \lim_1(f, \mathcal{F}_1)$ . The theorem is a consequence of (99).
- (103) Suppose  $\lim_{\langle \mathcal{F}_1, \mathcal{F}_2 \rangle} f \neq \emptyset$  and for every element  $x$  of  $X_1$ ,  $\lim_{\mathcal{F}_2} \text{curry}(f, x) \neq \emptyset$  and for every element  $x$  of  $X_2$ ,  $\lim_{\mathcal{F}_1} \text{curry}'(f, x) \neq \emptyset$ . Then  $\lim_{\mathcal{F}_1} \lim_2(f, \mathcal{F}_2) = \lim_{\mathcal{F}_2} \lim_1(f, \mathcal{F}_1)$ . The theorem is a consequence of (102) and (101).

## REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Formalized Mathematics*, 1(1):41–46, 1990.
- [2] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Formalized Mathematics*, 6(1):93–107, 1997.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [4] Grzegorz Bancerek, Noboru Endou, and Yuji Sakai. On the characterizations of compactness. *Formalized Mathematics*, 9(4):733–738, 2001.
- [5] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Kornilowicz, Roman Matuszewski, Adam Naumowicz, Karol Pał, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [6] Nicolas Bourbaki. *Topologie générale: Chapitres 1 à 4*. Éléments de mathématique. Springer Science & Business Media, 2007.
- [7] Nicolas Bourbaki. *General Topology: Chapters 1–4*. Springer Science and Business Media, 2013.
- [8] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [9] Czesław Byliński. Functions and their basic properties. *Formalized Mathematics*, 1(1):55–65, 1990.
- [10] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [11] Czesław Byliński. Some basic properties of sets. *Formalized Mathematics*, 1(1):47–53, 1990.
- [12] Roland Coghetto. Convergent filter bases. *Formalized Mathematics*, 23(3):189–203, 2015. doi:10.1515/forma-2015-0016.
- [13] Roland Coghetto. Summable family in a commutative group. *Formalized Mathematics*, 23(4):279–288, 2015. doi:10.1515/forma-2015-0022.

- [14] Noboru Endou, Hiroyuki Okazaki, and Yasunari Shidama. Double sequences and limits. *Formalized Mathematics*, 21(3):163–170, 2013. doi:10.2478/forma-2013-0018.
- [15] Andrzej Owsiejczuk. Combinatorial Grassmannians. *Formalized Mathematics*, 15(2):27–33, 2007. doi:10.2478/v10037-007-0004-9.
- [16] Karol Pałk. Stirling numbers of the second kind. *Formalized Mathematics*, 13(2):337–345, 2005.
- [17] Claude Wagschal. *Topologie et analyse fonctionnelle*. Hermann, 1995.
- [18] Claude Wagschal. *Topologie: Exercices et problèmes corrigés*. Hermann, 1995.
- [19] Edmund Woronowicz. Relations and their basic properties. *Formalized Mathematics*, 1(1):73–83, 1990.

*Received June 30, 2016*

---



The English version of this volume of *Formalized Mathematics* was financed under agreement 548/P-DUN/2016 with the funds from the Polish Minister of Science and Higher Education for the dissemination of science.