


Formal Development of Rough Inclusion Functions

Adam Grabowski 
Institute of Informatics
University of Białystok
Poland

Summary. Rough sets, developed by Pawlak [15], are important tool to describe situation of incomplete or partially unknown information. In this article, continuing the formalization of rough sets [12], we give the formal characterization of three rough inclusion functions (RIFs). We start with the standard one, $\kappa^{\mathcal{L}}$, connected with Łukasiewicz [14], and extend this research for two additional RIFs: κ_1 , and κ_2 , following a paper by Gomolińska [4], [3]. We also define q-RIFs and weak q-RIFs [2]. The paper establishes a formal counterpart of [7] and makes a preliminary step towards rough mereology [16], [17] in Mizar [13].

MSC: 03E70 68T99 03B35

Keywords: rough set; rough inclusion; approximation space

MML identifier: ROUGHIF1, version: 8.1.09 5.59.1363

0. INTRODUCTION

In the paper, continuing our development of rough sets, we define three closely related rough inclusion functions (RIFs).

Until now, most of the Mizar formalization of rough sets [5], [8] was done by means of the notion of a generalized approximation space understood as a pair $\langle U, \rho \rangle$, where ρ is an indiscernibility relation defined on the universe U . This viewpoint, based on tolerances instead of equivalence relations, was studied by Skowron and Stepaniuk [18], then, in a general form, by Zhu [19], among many others, and the Mizar counterpart of it is included in [9] and [10].

In the alternative approach, used by Gomolińska [3], approximation spaces are treated as triples of the form

$$\mathcal{A} = (U, I, \kappa),$$

where U is a non-empty set called the universe, $I : U \mapsto \wp U$ is an uncertainty mapping, and $\kappa : \wp U \times \wp U \mapsto [0, 1]$ is a rough inclusion function. The formalization of uncertainty mappings was discussed in [12], and this article tries to define the missing part of the above definition, with future possibility of merging approaches via theory merging mechanism [6], avoiding duplication as much as we can [11].

We start with some preliminaries, which cover gaps in the existing state of the Mizar Mathematical Library. Section 2 deals with the standard rough inclusion function, which appears in some form in the research by Jan Łukasiewicz [14], obviously without any reference for rough sets. This pretty general Mizar functor $\kappa^{\mathcal{L}}$ is defined as follows:

$$\kappa^{\mathcal{L}}(X, Y) = \begin{cases} \frac{|X \cap Y|}{|X|}, & \text{if } X \neq \emptyset \\ 1, & \text{otherwise} \end{cases}$$

For a given universe U , rough inclusion functions (RIFs for short) are the mappings κ from $\wp U \times \wp U$ into unit interval which satisfy two properties:

$$\text{rif}_1(\kappa) \Leftrightarrow \forall X, Y \subseteq U (\kappa(X, Y) = 1 \Leftrightarrow X \subseteq Y)$$

$$\text{rif}_2(\kappa) \Leftrightarrow \forall X, Y, Z \subseteq U (Y \subseteq Z \Rightarrow \kappa(X, Y) \leq \kappa(X, Z))$$

This is discussed in Sect. 3; corresponding Mizar modes **RIF** and **preRIF** are also introduced.

Besides $\kappa^{\mathcal{L}}$, there are two relatively well-known RIFs:

$$\kappa_1(X, Y) = \begin{cases} \frac{|Y|}{|X \cup Y|}, & \text{if } X \cup Y \neq \emptyset \\ 1, & \text{otherwise} \end{cases}$$

$$\kappa_2(X, Y) = \frac{|(U - X) \cup Y|}{|U|}.$$

Section 4 contains their definitions, both of the form of Mizar functors, and as set-theoretic functions.

It should be mentioned that in this Mizar translation (and also in the source code), predicative form of the properties of RIFs, as, for example, $\text{rif}_1(\kappa)$ is replaced by the phrase “ κ satisfies (RIF₁)” (and for others, respectively). In Sect. 5 we formulate some additional characteristic properties of rough inclusions; in Sect. 6 we show that, under the assumption that rif_1 holds, rif_2 can be

replaced by rif_2^* . We introduce also some weakened versions of rough inclusions: quasi-RIF and weak quasi-RIF.

All three considered RIFs $(\kappa^\mathcal{L}, \kappa_1, \kappa_2)$ are distinct. Gomolińska takes $U = \{0, 1, 2, \dots, 9\}$, $X = \{0, \dots, 4\}$, $Y = \{2, \dots, 6\}$. Then $\kappa^\mathcal{L}(X, Y) = 3/5$, $\kappa_1(X, Y) = 5/7$, and $\kappa_2(X, Y) = 4/5$. In Sect. 9, we constructed an example, which in Mizar is the functor `ExampleRIFSpace`, claiming that $U = \{1, 2, 3, 4, 5\}$, $X = \{1, 2\}$, $Y = \{2, 3, 4\}$ with $\kappa^\mathcal{L}(X, Y) = 1/2$, $\kappa_1(X, Y) = 3/4$, and $\kappa_2(X, Y) = 4/5$. Obviously, the indiscernibility relation does not matter, and so we took the identity as the simplest. The proofs, based on our specific example, are significantly shorter than those proposed by Gomolińska.

In the final section, we formalized two theorems from another Gomolińska's paper [1], which was already translated into Mizar [12], but without the notion of RIF; now we can fill this gap.

1. PRELIMINARIES

Now we state the propositions:

- (1) Let us consider real numbers a, b, c . If $b > 0$ and $a \leq b$ and $c \geq 0$, then $\frac{a}{b} \leq \frac{a+c}{b+c}$.

Observe that there exists an approximation space which is strict and finite.

Let R be a finite 1-sorted structure. One can check that every subset of R is finite.

From now on R denotes a 1-sorted structure and X, Y denote subsets of R .

- (2) $X \subseteq Y$ if and only if $X^c \cup Y = \Omega_R$.

PROOF: If $X \subseteq Y$, then $X^c \cup Y = \Omega_R$. \square

From now on R denotes a finite 1-sorted structure and X, Y denote subsets of R . Now we state the propositions:

- (3) $\overline{X \cup Y} = \overline{Y}$ if and only if $X \subseteq Y$.
- (4) If $\overline{X^c \cup Y} = \overline{\Omega_R}$, then $X^c \cup Y = \Omega_R$.

Let R be a non empty 1-sorted structure and X be a subset of R . Note that $\Omega_R \cup X$ reduces to Ω_R and $\Omega_R \cap X$ reduces to X .

2. STANDARD ROUGH INCLUSION FUNCTION

From now on R denotes a finite approximation space and X, Y, Z, W denote subsets of R .

Let R be a finite approximation space and X, Y be subsets of R . The functor $\kappa^\mathcal{L}(X, Y)$ yielding an element of $[0, 1]$ is defined by the term

$$(Def. 1) \quad \begin{cases} \frac{\overline{\overline{X \cap Y}}}{\overline{X}}, & \text{if } X \neq \emptyset, \\ 1, & \text{otherwise.} \end{cases}$$

Now we state the propositions:

- (5) $\kappa^{\mathcal{L}}(\emptyset_R, X) = 1$.
- (6) $\kappa^{\mathcal{L}}(X, Y) = 1$ if and only if $X \subseteq Y$.
- (7) If $Y \subseteq Z$, then $\kappa^{\mathcal{L}}(X, Y) \leq \kappa^{\mathcal{L}}(X, Z)$.
- (8) If $Z \subseteq Y \subseteq X$, then $\kappa^{\mathcal{L}}(X, Z) \leq \kappa^{\mathcal{L}}(Y, Z)$.
- (9) $\kappa^{\mathcal{L}}(X, Y \cup Z) \leq \kappa^{\mathcal{L}}(X, Y) + \kappa^{\mathcal{L}}(X, Z)$.
- (10) If $X \neq \emptyset$ and Y misses Z , then $\kappa^{\mathcal{L}}(X, Y \cup Z) = \kappa^{\mathcal{L}}(X, Y) + \kappa^{\mathcal{L}}(X, Z)$.

3. ROUGH INCLUSION FUNCTIONS

Let R be a 1-sorted structure.

A pre-rough inclusion function of R is a function from $2^{(\text{the carrier of } R)} \times 2^{(\text{the carrier of } R)}$ into $[0, 1]$.

A preRIF of R is a pre-rough inclusion function of R .

The scheme *BinOpEq* deals with a non empty 1-sorted structure \mathcal{R} and a binary functor \mathcal{F} yielding an element of $[0, 1]$ and states that

- (Sch. 1) For every preRIFs f_1, f_2 of \mathcal{R} such that for every subsets x, y of \mathcal{R} , $f_1(x, y) = \mathcal{F}(x, y)$ and for every subsets x, y of \mathcal{R} , $f_2(x, y) = \mathcal{F}(x, y)$ holds $f_1 = f_2$.

Let R be a finite approximation space. The functor $\kappa^{\mathcal{L}}(R)$ yielding a preRIF of R is defined by

- (Def. 2) for every subsets x, y of R , $it(x, y) = \kappa^{\mathcal{L}}(x, y)$.

4. DEFINING TWO NEW RIFs

Let R be a finite approximation space and X, Y be subsets of R . The functor $\kappa_1(X, Y)$ yielding an element of $[0, 1]$ is defined by the term

$$(Def. 3) \quad \begin{cases} \frac{\overline{\overline{Y}}}{\overline{\overline{X \cup Y}}}, & \text{if } X \cup Y \neq \emptyset, \\ 1, & \text{otherwise.} \end{cases}$$

The functor $\kappa_2(X, Y)$ yielding an element of $[0, 1]$ is defined by the term

$$(Def. 4) \quad \frac{\overline{\overline{X \cup Y}}}{\Omega_R}.$$

The functor $\kappa_1(R)$ yielding a preRIF of R is defined by

- (Def. 5) for every subsets x, y of R , $it(x, y) = \kappa_1(x, y)$.

The functor $\kappa_2(R)$ yielding a preRIF of R is defined by

(Def. 6) for every subsets x, y of R , $it(x, y) = \kappa_2(x, y)$.

Now we state the propositions:

- (11) $\kappa_1(X, Y) = 1$ if and only if $X \subseteq Y$. The theorem is a consequence of (3).
- (12) $\kappa_2(X, Y) = 1$ if and only if $X \subseteq Y$. The theorem is a consequence of (2) and (4).
- (13) If $\kappa_1(X, Y) = 0$, then $Y = \emptyset$.
- (14) If $X \neq \emptyset$, then $\kappa_1(X, Y) = 0$ iff $Y = \emptyset$. The theorem is a consequence of (13).
- (15) $\kappa_2(X, Y) = 0$ if and only if $X = \Omega_R$ and $Y = \emptyset$.
- (16) If $Y \subseteq Z$, then $\kappa_1(X, Y) \leq \kappa_1(X, Z)$. The theorem is a consequence of (1).
- (17) If $Y \subseteq Z$, then $\kappa_2(X, Y) \leq \kappa_2(X, Z)$.
- (18) $\kappa_1(\emptyset_R, X) = 1$. The theorem is a consequence of (11).
- (19) $\kappa_2(\emptyset_R, X) = 1$. The theorem is a consequence of (12).

5. CHARACTERISTIC PROPERTIES OF ROUGH INCLUSIONS

Let R be a non empty relational structure and κ be a preRIF of R . We say that κ satisfies (RIF₁) if and only if

(Def. 7) for every subsets X, Y of R , $\kappa(X, Y) = 1$ iff $X \subseteq Y$.

We say that κ satisfies (RIF₂) if and only if

(Def. 8) for every subsets X, Y, Z of R such that $Y \subseteq Z$ holds $\kappa(X, Y) \leq \kappa(X, Z)$.

We say that κ satisfies (RIF₃) if and only if

(Def. 9) for every subset X of R such that $X \neq \emptyset$ holds $\kappa(X, \emptyset_R) = 0$.

We say that κ satisfies (RIF₄) if and only if

(Def. 10) for every subsets X, Y of R such that $\kappa(X, Y) = 0$ holds X misses Y .

We say that κ satisfies (RIF₀) if and only if

(Def. 11) for every subsets X, Y of R such that $X \subseteq Y$ holds $\kappa(X, Y) = 1$.

We say that κ satisfies (RIF₀⁻¹) if and only if

(Def. 12) for every subsets X, Y of R such that $\kappa(X, Y) = 1$ holds $X \subseteq Y$.

We say that κ satisfies (RIF₂^{*}) if and only if

(Def. 13) for every subsets X, Y, Z of R such that $\kappa(Y, Z) = 1$ holds $\kappa(X, Y) \leq \kappa(X, Z)$.

Observe that every preRIF of R which satisfies (RIF_1) satisfies also (RIF_0) and (RIF_0^{-1}) and every preRIF of R which satisfies (RIF_0) and (RIF_0^{-1}) satisfies also (RIF_1) .

Let R be a finite approximation space. One can check that $\kappa^{\mathcal{L}}(R)$ satisfies (RIF_1) and $\kappa^{\mathcal{L}}(R)$ satisfies (RIF_2) and $\kappa_1(R)$ satisfies (RIF_1) and $\kappa_1(R)$ satisfies (RIF_2) and $\kappa_2(R)$ satisfies (RIF_1) and $\kappa_2(R)$ satisfies (RIF_2) .

Let us consider R . Note that there exists a preRIF of R which satisfies (RIF_1) and (RIF_2) .

6. ON THE CONNECTIONS BETWEEN POSTULATES

Now we state the proposition:

- (20) Let us consider preRIF κ of R satisfying (RIF_1) . Then κ satisfies (RIF_2) if and only if κ satisfies (RIF_2^*) .

Let us consider R . Let us observe that every preRIF of R satisfying (RIF_1) which satisfies (RIF_2) satisfies also (RIF_2^*) and every preRIF of R satisfying (RIF_1) which satisfies (RIF_2^*) satisfies also (RIF_2) and $\kappa^{\mathcal{L}}(R)$ satisfies (RIF_0) and (RIF_2^*) and there exists a pre-rough inclusion function of R which satisfies (RIF_0) , (RIF_1) , (RIF_2) , and (RIF_2^*) .

A rough inclusion function of R is pre-rough inclusion function of R satisfying (RIF_1) and (RIF_2) .

A quasi-rough inclusion function of R is preRIF of R satisfying (RIF_0) and (RIF_2^*) .

A weak quasi-rough inclusion function of R is preRIF of R satisfying (RIF_0) and (RIF_2) .

A RIF of R is a rough inclusion function of R .

A q-RIF of R is a quasi-rough inclusion function of R .

A weak q-RIF of R is a weak quasi-rough inclusion function of R .

7. FORMALIZATION OF PROPOSITION 2 [3]

Now we state the propositions:

- (21) If $X \neq \emptyset$ and $Z \cup W = \Omega_R$ and Z misses W , then $\kappa^{\mathcal{L}}(X, Z) + \kappa^{\mathcal{L}}(X, W) = 1$.
- (22) If $\kappa^{\mathcal{L}}(X, Y) = 0$, then X misses Y .
- (23) If $X \neq \emptyset$, then $\kappa^{\mathcal{L}}(X, Y) = 0$ iff X misses Y . The theorem is a consequence of (22).
- (24) If $X \neq \emptyset$, then $\kappa^{\mathcal{L}}(X, \emptyset_R) = 0$.

Now we state the propositions:

- (25) If $X \neq \emptyset$ and X misses Y , then $\kappa^{\mathcal{L}}(X, Z \setminus Y) = \kappa^{\mathcal{L}}(X, Z \cup Y) = \kappa^{\mathcal{L}}(X, Z)$.
The theorem is a consequence of (23), (10), (7), and (9).
- (26) If Z misses W , then $\kappa^{\mathcal{L}}(Y \cup Z, W) \leq \kappa^{\mathcal{L}}(Y, W) \leq \kappa^{\mathcal{L}}(Y \setminus Z, W)$.
- (27) If Z misses Y and $Z \subseteq W$, then $\kappa^{\mathcal{L}}(Y \setminus Z, W) \leq \kappa^{\mathcal{L}}(Y, W) \leq \kappa^{\mathcal{L}}(Y \cup Z, W)$.

8. FORMALIZATION OF PROPOSITION 4 [3]

Let us consider R . Let X be a non empty subset of R . Let us note that $\kappa^{\mathcal{L}}(X, \emptyset_R)$ is empty.

Now we state the propositions:

- (28) If $\kappa_1(X, Y) = 0$, then X misses Y . The theorem is a consequence of (14).
- (29) If $\kappa_2(X, Y) = 0$, then X misses Y . The theorem is a consequence of (15).

Let us consider R . Observe that $\kappa^{\mathcal{L}}(R)$ satisfies (RIF₄) and $\kappa_1(R)$ satisfies (RIF₄) and $\kappa_2(R)$ satisfies (RIF₄).

- (30) $\kappa^{\mathcal{L}}(X, Y) \leq \kappa_1(X, Y) \leq \kappa_2(X, Y)$. The theorem is a consequence of (1), (18), and (19).
- (31) $\kappa_1(X, Y) = \kappa^{\mathcal{L}}(X \cup Y, Y)$. The theorem is a consequence of (6) and (11).
- (32) $\kappa_2(X, Y) = \kappa^{\mathcal{L}}(\Omega_R, X^c \cup Y) = \kappa^{\mathcal{L}}(\Omega_R, X^c) + \kappa^{\mathcal{L}}(\Omega_R, X \cap Y)$.
- (33) $\kappa^{\mathcal{L}}(X, Y) = \kappa^{\mathcal{L}}(X, X \cap Y) = \kappa_1(X, X \cap Y) = \kappa_1(X \setminus Y, X \cap Y)$.
- (34) If $X \cup Y = \Omega_R$, then $\kappa_1(X, Y) = \kappa_2(X, Y)$. The theorem is a consequence of (2).
- (35) If $X \neq \emptyset$, then $1 - \kappa^{\mathcal{L}}(X, Y) = \kappa^{\mathcal{L}}(X, Y^c)$. The theorem is a consequence of (10) and (6).

9. CONCRETE EXAMPLE

Let X be a set. The functor $\text{DiscreteApproxSpace}(X)$ yielding a strict relational structure is defined by the term

(Def. 14) $\langle X, \text{id}_X \rangle$.

Let us note that $\text{DiscreteApproxSpace}(X)$ has equivalence relation.

Let X be a non empty set. Observe that $\text{DiscreteApproxSpace}(X)$ is non empty.

Let X be a finite set. Let us observe that $\text{DiscreteApproxSpace}(X)$ is finite.

The functor ExampleRIFSpace yielding a strict, finite approximation space is defined by the term

(Def. 15) $\text{DiscreteApproxSpace}(\{1, 2, 3, 4, 5\})$.

Now we state the propositions:

- (36) Let us consider subsets X, Y of ExampleRIFSpace . Suppose $X = \{1, 2\}$ and $Y = \{2, 3, 4\}$. Then $\kappa^{\mathcal{L}}(X, Y) \neq \kappa^{\mathcal{L}}(Y, X)$.
- (37) Let us consider subsets X, Y, U of ExampleRIFSpace . Suppose $X = \{1, 2\}$ and $Y = \{1, 2, 3\}$ and $U = \{2, 4, 5\}$. Then $\kappa^{\mathcal{L}}(X, U) \not\leq \kappa^{\mathcal{L}}(Y, U)$.
- (38) Let us consider subsets X, Y of ExampleRIFSpace . Suppose $X = \{1, 2\}$ and $Y = \{2, 3, 4\}$. Then $\kappa^{\mathcal{L}}(X, Y)$, $\kappa_1(X, Y)$, $\kappa_2(X, Y)$ are mutually different.

10. CONTINUING FORMALIZATION OF THEOREM 4.1 [1]

Let us consider a finite approximation space R , an element u of R , and subsets x, y of R . Now we state the propositions:

- (39) If $u \in (f_1(R))(x)$ and $(I_R)(u) = y$, then $\kappa^{\mathcal{L}}(y, x) > 0$. The theorem is a consequence of (22).
- (40) If $u \in (\text{Flip } f_1(R))(x)$ and $(I_R)(u) = y$, then $\kappa^{\mathcal{L}}(y, x) = 1$. The theorem is a consequence of (6).

REFERENCES

- [1] Anna Gomolińska. A comparative study of some generalized rough approximations. *Fundamenta Informaticae*, 51:103–119, 2002.
- [2] Anna Gomolińska. Rough approximation based on weak q-RIFs. In James F. Peters, Andrzej Skowron, Marcin Wolski, Mihir K. Chakraborty, and Wei-Zhi Wu, editors, *Transactions on Rough Sets X*, volume 5656 of *Lecture Notes in Computer Science*, pages 117–135, Berlin, Heidelberg, 2009. Springer. ISBN 978-3-642-03281-3. doi:10.1007/978-3-642-03281-3_4.
- [3] Anna Gomolińska. On three closely related rough inclusion functions. In Marzena Kryszkiewicz, James F. Peters, Henryk Rybiński, and Andrzej Skowron, editors, *Rough Sets and Intelligent Systems Paradigms*, volume 4585 of *Lecture Notes in Computer Science*, pages 142–151, Berlin, Heidelberg, 2007. Springer. doi:10.1007/978-3-540-73451-2_16.
- [4] Anna Gomolińska. On certain rough inclusion functions. In James F. Peters, Andrzej Skowron, and Henryk Rybiński, editors, *Transactions on Rough Sets IX*, volume 5390 of *Lecture Notes in Computer Science*, pages 35–55. Springer Berlin Heidelberg, 2008. doi:10.1007/978-3-540-89876-4_3.
- [5] Adam Grabowski. On the computer-assisted reasoning about rough sets. In B. Dunin-Kępicz, A. Jankowski, A. Skowron, and M. Szczuka, editors, *International Workshop on Monitoring, Security, and Rescue Techniques in Multiagent Systems Location*, volume 28 of *Advances in Soft Computing*, pages 215–226, Berlin, Heidelberg, 2005. Springer-Verlag. doi:10.1007/3-540-32370-8_15.
- [6] Adam Grabowski. Efficient rough set theory merging. *Fundamenta Informaticae*, 135(4): 371–385, 2014. doi:10.3233/FI-2014-1129.
- [7] Adam Grabowski. Building a framework of rough inclusion functions by means of computerized proof assistant. In Tamás Mihálydeák, Fan Min, Guoyin Wang, Mohua Banerjee, Ivo Düntsch, Zbigniew Suraj, and Davide Ciucci, editors, *Rough Sets*, volume 11499 of *Lecture Notes in Computer Science*, pages 225–238, Cham, 2019. Springer International Publishing. ISBN 978-3-030-22815-6. doi:10.1007/978-3-030-22815-6_18.

- [8] Adam Grabowski. Lattice theory for rough sets – a case study with Mizar. *Fundamenta Informaticae*, 147(2–3):223–240, 2016. doi:10.3233/FI-2016-1406.
- [9] Adam Grabowski. Relational formal characterization of rough sets. *Formalized Mathematics*, 21(1):55–64, 2013. doi:10.2478/forma-2013-0006.
- [10] Adam Grabowski. Binary relations-based rough sets – an automated approach. *Formalized Mathematics*, 24(2):143–155, 2016. doi:10.1515/forma-2016-0011.
- [11] Adam Grabowski and Christoph Schwarzweiler. On duplication in mathematical repositories. In Serge Autexier, Jacques Calmet, David Delahaye, Patrick D. F. Ion, Laurence Rideau, Renaud Rioboo, and Alan P. Sexton, editors, *Intelligent Computer Mathematics, 10th International Conference, AISC 2010, 17th Symposium, Calculemus 2010, and 9th International Conference, MKM 2010, Paris, France, July 5–10, 2010. Proceedings*, volume 6167 of *Lecture Notes in Computer Science*, pages 300–314. Springer, 2010. doi:10.1007/978-3-642-14128-7_26.
- [12] Adam Grabowski and Michał Sielwiesiuk. Formalizing two generalized approximation operators. *Formalized Mathematics*, 26(2):183–191, 2018. doi:10.2478/forma-2018-0016.
- [13] Adam Grabowski, Artur Kornilowicz, and Adam Naumowicz. Four decades of Mizar. *Journal of Automated Reasoning*, 55(3):191–198, 2015. doi:10.1007/s10817-015-9345-1.
- [14] Jan Łukasiewicz. Die logischen Grundlagen der Wahrscheinlichkeitsrechnung. In L. Borkowski, editor, *Jan Łukasiewicz – Selected Works*, pages 16–63. North Holland, Polish Scientific Publ., Amsterdam London Warsaw, 1970. First published in Kraków, 1913.
- [15] Zdzisław Pawlak. Rough sets. *International Journal of Parallel Programming*, 11:341–356, 1982. doi:10.1007/BF01001956.
- [16] Lech Polkowski. Rough mereology. In *Approximate Reasoning by Parts*, volume 20 of *Intelligent Systems Reference Library*, pages 229–257, Berlin, Heidelberg, 2011. Springer. ISBN 978-3-642-22279-5. doi:10.1007/978-3-642-22279-5_6.
- [17] Lech Polkowski and Andrzej Skowron. Rough mereology: A new paradigm for approximate reasoning. *International Journal of Approximate Reasoning*, 15(4):333–365, 1996. doi:10.1016/S0888-613X(96)00072-2.
- [18] Andrzej Skowron and Jarosław Stepaniuk. Tolerance approximation spaces. *Fundamenta Informaticae*, 27(2/3):245–253, 1996. doi:10.3233/FI-1996-272311.
- [19] William Zhu. Generalized rough sets based on relations. *Information Sciences*, 177:4997–5011, 2007.

Accepted August 29, 2019
